

Biogeochemical Cycling

Raina M. Maier

16.1 Introduction	16.3.3 Ammonia Assimilation (Immobilization) and Ammonification (Mineralization)	16.4.3 Sulfur Oxidation
16.1.1 Biogeochemical Cycles		16.4.4 Sulfur Reduction
16.1.2 Gaia Hypothesis		16.5 Iron Cycle
16.2 Carbon Cycle	16.3.4 Ammonium Oxidation	16.5.1 Iron Reservoirs
16.2.1 Carbon Reservoirs	16.3.5 Nitrate Reduction	16.5.2 Iron in Soils and Sediments
16.2.2 Carbon Fixation and Energy Flow	16.4 Sulfur Cycle	16.5.3 Iron in Marine Environments
16.2.3 Carbon Respiration	16.4.1 Sulfur Reservoirs	16.5.4 Iron Oxidation
16.3 Nitrogen Cycle	16.4.2 Assimilatory Sulfate Reduction and Sulfur Mineralization	16.5.5 Iron Reduction
16.3.1 Nitrogen Reservoirs		Questions and Problems
16.3.2 Nitrogen Fixation		References and Recommended Reading

16.1 INTRODUCTION

16.1.1 Biogeochemical Cycles

Carbon dioxide concentrations in the atmosphere exceeded 400 parts per million (ppm) for the first time in May 2013, increasing from 315 ppm in 1958 (when the first accurate measurements were made), reports the U.S. National Oceanic and Atmospheric Administration. What has driven this increase? Carbon is cycled between organic forms such as sugar or other cellular building blocks and inorganic forms such as carbon dioxide. Vast amounts of organic matter are photosynthetically produced on Earth each year utilizing carbon dioxide from the atmosphere. Most of this material is consumed and degraded but a part of it, over the millennia, has been stored in permafrost, peat bogs and as fossil fuels. A delicate global balance of organic and inorganic carbon has been maintained largely driven by microbial activity. Human use of stored organic carbon (fossil fuels and peat) and recent warming of permafrost have upset this balance in favor of the release of carbon dioxide to the

atmosphere. Most scientists agree that this is having a major impact on global warming, and is an excellent example of a major perturbation of the carbon cycle—one that is occurring during our lifetimes. Cycling between organic and inorganic forms is not limited to carbon. All of the major elements found in biological organisms (see [Table 16.1](#)), as well as some of the minor and trace elements, are similarly cycled in predictable and definable ways.

Taken together, the various element cycles are called the **biogeochemical cycles**. Understanding these cycles allows scientists to understand and predict the development of microbial communities and activities in the environment. There are many activities that can be harnessed in a beneficial way, such as for remediation of organic and metal pollutants, or for recovery of precious metals such as copper or uranium from low-grade ores. There are detrimental aspects of the cycles that can cause global environmental problems, for example the formation of acid rain and acid mine drainage, metal corrosion processes and formation of nitrous oxide, which can deplete Earth's ozone layer (see Chapter 31). As these examples

illustrate, the microbial activities that drive biogeochemical cycles are highly relevant to the field of environmental microbiology. Thus, the knowledge of these cycles is increasingly critical as the human population continues to grow, and the impact of human activity on Earth's environment becomes more significant. In this chapter,

the biogeochemical cycles pertaining to carbon, nitrogen, sulfur and iron are delineated. Although the discussion will be limited to these four cycles, it should be noted that there are a number of other cycles—the phosphorus cycle, the manganese cycle, the calcium cycle and more (Dobrovolsky, 1994).

TABLE 16.1 Chemical Composition of an *E. coli* Cell

Elemental Breakdown	% Dry Mass of an <i>E. coli</i> Cell
Major elements	
Carbon	50
Oxygen	20
Hydrogen	8
Nitrogen	14
Sulfur	1
Phosphorus	3
Minor elements	
Potassium	2
Calcium	0.05
Magnesium	0.05
Chlorine	0.05
Iron	0.2
Trace elements	
Manganese	all trace elements
Molybdenum	combined comprise 0.3%
Cobalt	of dry weight of cell
Copper	
Zinc	

Adapted from Neidhardt et al. (1990).

16.1.2 Gaia Hypothesis

In the early 1970s, James Lovelock theorized that Earth behaves like a superorganism, and this concept developed into what is now known as the **Gaia hypothesis**. To quote Lovelock (1995), “Living organisms and their material environment are tightly coupled. The coupled system is a superorganism, and as it evolves there emerges a new property, the ability to self-regulate climate and chemistry.” The basic tenet of this hypothesis is that Earth's physicochemical properties are self-regulated so that they are maintained in a favorable range for life. As evidence for this, consider that the sun has heated up by 30% during the past 4–5 billion years. Given Earth's original carbon dioxide-rich atmosphere, the average surface temperature of a lifeless Earth today would be approximately 290°C (Table 16.2). In fact, when one compares Earth's present-day atmosphere with the atmospheres found on our nearest neighbors Venus and Mars, one can see that something has drastically affected the development of Earth's atmosphere. According to the Gaia hypothesis, this is the development and continued presence of life. Microbial activity, and later the appearance of plants, have changed the original heat-trapping carbon dioxide-rich atmosphere to the present oxidizing, carbon dioxide-poor atmosphere. This has allowed Earth to maintain an average surface temperature of 13°C, which is favorable to the life that exists on Earth.

How do biogeochemical activities relate to the Gaia hypothesis? These biological activities have driven

TABLE 16.2 Atmosphere and Temperatures Found on Venus, Mars and Earth

Gas	Planet			
	Venus	Mars	Earth Without Life	Earth With Life
Carbon dioxide	96.5%	95%	98%	0.03%
Nitrogen	3.5%	2.7%	1.9%	9%
Oxygen	trace	0.13%	0.0	21%
Argon	70 ppm	1.6%	0.1%	1%
Methane	0.0	0.0	0.0	1.7 ppm
Surface temperature (°C)	459	−53	290 ± 50	13

Adapted from Lovelock (1995).



FIGURE 16.1 An example of a living stromatolite (left) and a stromatolite fossil (right). From (left) Reynolds (1999) and (right) Farabee (2008).

the response to the slow warming of the sun resulting in the major atmospheric changes that have occurred over the last 4–5 billion years. When Earth was formed 4–5 billion years ago, a reducing (anaerobic) atmosphere existed. The initial reactions that mediated the formation of organic carbon were abiotic, driven by large influxes of ultraviolet (UV) light. The resulting reservoir of organic matter was utilized by early anaerobic heterotrophic organisms. This was followed by the development of the ability of microbes to fix carbon dioxide photosynthetically. Evidence from **stromatolite** fossils suggests that the ability to photosynthesize was developed at least 3.5 billion years ago. Stromatolites are fossilized laminated structures that have been found in Africa and Australia (Figure 16.1). Although a hotly debated topic, there is evidence that these structures were formed by photosynthetic microorganisms (first anaerobic, then cyanobacterial) that grew in mats and entrapped or precipitated inorganic material as they grew (Bosak *et al.*, 2007).

The evolution of photosynthetic organisms tapped into an unlimited source of energy, the sun, and provided a mechanism for carbon recycling, i.e., the first carbon cycle (Figure 16.2). This first carbon cycle was maintained for approximately 1.5 billion years. Geologic evidence then suggests that approximately 2 billion years ago, photosynthetic microorganisms developed the ability to produce oxygen. This allowed oxygen to accumulate in the atmosphere, resulting, in time, in a change from reducing to oxidizing conditions. Further, oxygen accumulation in the atmosphere created an ozone layer, which reduced the influx of harmful UV radiation, allowing the development of higher forms of life to begin.

At the same time that the carbon cycle evolved, the nitrogen cycle emerged because nitrogen was a limiting element for microbial growth. Although molecular nitrogen was abundant in the atmosphere, microbial cells




FIGURE 16.2 The carbon cycle is dependent on autotrophic organisms that fix carbon dioxide into organic carbon and heterotrophic organisms that convert organic carbon to carbon dioxide during respiration.

could not directly utilize nitrogen as N_2 gas. Cells require organic nitrogen compounds or reduced inorganic forms of nitrogen for growth. Therefore, under the reducing conditions found on early Earth, some organisms developed a mechanism for fixing nitrogen using the enzyme nitrogenase. Nitrogen fixation remains an important microbiological process, and to this day the majority of nitrogenase enzymes are totally inhibited in the presence of oxygen.

When considered over this geologic time scale of several billion years, it is apparent that biogeochemical activities have been unidirectional. This means that the predominant microbial activities on Earth have evolved over this long period of time to produce changes, and to respond to changes that have occurred in the atmosphere, i.e., the appearance of oxygen and the decrease in carbon dioxide content. Presumably these changes will continue to occur, but they occur so slowly that we do not have the capacity to observe them.

TABLE 16.3 Global Carbon Reservoirs

Carbon Reservoir	Metric Tons Carbon	Actively Cycled
Atmosphere		
CO ₂	6.7×10^{11}	Yes
Ocean		
Biomass	4.0×10^9	No
Carbonates	3.8×10^{13}	No
Dissolved and particulate organics	2.1×10^{12}	Yes
Land		
Biota	5.0×10^{11}	Yes
Humus	1.2×10^{12}	Yes
Fossil fuel	1.0×10^{13}	Yes
Earth's crust ^a	1.2×10^{17}	No

Data from Dobrovolsky (1994).

^aThis reservoir includes the entire lithosphere found in either terrestrial or ocean environments.

One can also consider biogeochemical activities on a more contemporary time scale, that of tens to hundreds of years. On this much shorter time scale, biogeochemical activities are regular and cyclic in nature, and it is these activities that are addressed in this chapter. On the one hand, the presumption that Earth is a superorganism that can respond to drastic environmental changes is heartening when one considers that human activity is causing unexpected changes in the atmosphere, such as ozone depletion and buildup of carbon dioxide. However, it is important to point out that the response of a superorganism is necessarily slow (thousands to millions of years), and as residents of Earth we must be sure not to overtax Earth's ability to respond to change by artificially changing the environment in a much shorter time frame.

16.2 CARBON CYCLE

16.2.1 Carbon Reservoirs

A **reservoir** is a sink or source of an element such as carbon. There are various global reservoirs of carbon, some of which are immense in size and some of which are relatively small (Table 16.3). The largest carbon reservoir is carbonate rock found in Earth's crust. This reservoir is four orders of magnitude larger than the carbonate reservoir found in the ocean, and six orders of magnitude larger than the carbon reservoir found as carbon dioxide in the atmosphere. If one considers these three reservoirs,

TABLE 16.4 Net Carbon Flux between Selected Carbon Reservoirs

Carbon Source	Flux (Metric Tons Carbon/Year)
Release by fossil fuel combustion	7×10^9
Land clearing	3×10^9
Forest harvest and decay	6×10^9
Forest regrowth	-4×10^9
Net uptake by oceans (diffusion)	-3×10^9
Annual flux	9×10^9

Adapted from Atlas and Bartha (1993).

it is obvious that the carbon most available for photosynthesis, which requires carbon dioxide, is in the smallest of the reservoirs, the atmosphere. Therefore, it is the smallest reservoir that is most actively cycled. It is small, actively cycled reservoirs such as atmospheric carbon dioxide that are subject to perturbation from human activity. In fact, since global industrialization began in the late 1800s, humans have affected several of the smaller carbon reservoirs. Utilization of fossil fuels (an example of a small, inactive carbon reservoir) and deforestation (an example of a small, active carbon reservoir) are two activities that have reduced the amount of fixed organic carbon in these reservoirs, and added to the atmospheric carbon dioxide reservoir (Table 16.4).

The increase in atmospheric carbon dioxide has not been as great as expected. This is because the reservoir of carbonate found in the ocean acts as a buffer between the atmospheric and sediment carbon reservoirs through the equilibrium equation shown below:



Thus, some of the excess carbon dioxide that has been released has been absorbed by the oceans. However, there has still been a net efflux of carbon dioxide into the atmosphere of approximately 7×10^9 metric tons/year. The problem with this imbalance is that because atmospheric carbon dioxide is a small carbon reservoir, the result of a continued net efflux over the past 100 years or so has been a 28% increase in atmospheric carbon dioxide from 0.026 to 0.033%. A consequence of the increase in atmospheric carbon dioxide is that it contributes to global warming through the **greenhouse effect** (see also Chapter 31). The greenhouse effect is caused by gases in the atmosphere that trap heat from the sun and cause Earth to warm up. This effect is not solely due to carbon




FIGURE 16.3 Diagram of the efficiency of sunlight energy flow from primary producers to consumers.

dioxide; other gases such as **methane**, **chlorofluorocarbons (CFCs)** and **nitrous oxide** all contribute to the problem.

16.2.2 Carbon Fixation and Energy Flow

The ability to photosynthesize allows sunlight energy to be trapped and stored. In this process carbon dioxide is fixed into organic matter (Figure 16.2). Photosynthetic organisms, also called **primary producers**, include plants and microorganisms such as algae, cyanobacteria, some bacteria and some protozoa. As shown in Figure 16.3, the efficiency of sunlight trapping is very low; less than 0.1% of the sunlight energy that hits Earth is actually utilized. As the fixed sunlight energy moves up each level of the food chain, up to 90% or more of the trapped energy is lost through respiration. Despite this seemingly inefficient trapping, photoautotrophic primary producers support most of the considerable ecosystems found on Earth. Productivity varies widely among different ecosystems depending on the climate, the type of primary producer and whether the system is a managed one (Table 16.5). For example, one of the most productive natural areas is the coral reefs. Managed agricultural systems such as corn and sugarcane systems are also very productive, but it should be remembered that a significant amount of energy is put into these systems in terms of fertilizer addition and care. The open ocean has much lower productivity, but covers the majority of Earth's surface, and so is a major contributor to primary production. In fact, aquatic and terrestrial environments contribute almost equally to global primary production. Plants predominate in terrestrial environments, but with the exception of immediate coastal zones, microorganisms are responsible for most primary production in aquatic environments. It follows that microorganisms are responsible for approximately one-half of all primary production on Earth.

TABLE 16.5 Net Primary Productivity of Some Natural and Managed Ecosystems

Description of Ecosystem	Net Primary Productivity (g Dry Organic Matter/m ² /Year)
Tundra	400
Desert	200
Temperate grassland	Up to 1500
Temperate or deciduous forest	1200–1600
Tropical rain forest	Up to 2800
Cattail swamp	2500
Freshwater pond	950–1500
Open ocean	100
Coastal seawater	200
Upwelling area	600
Coral reef	4900
Corn field	1000–6000
Rice paddy	340–1200
Sugarcane field	Up to 9400

Adapted from Atlas and Bartha (1993).

16.2.3 Carbon Respiration

Carbon dioxide that is fixed into organic compounds as a result of photoautotrophic activity is available for consumption or respiration by animals and heterotrophic microorganisms. This is the second half of the carbon cycle shown in Figure 16.2. The end products of respiration are carbon dioxide and new cell mass. An interesting question to consider is the following: if respiration were to stop, how long would it take for photosynthesis to use up all of the carbon dioxide reservoir in the atmosphere? Based on estimates of global photosynthesis, it has been estimated that it would take 30 to 300 years. This illustrates the importance of both legs of the carbon cycle in maintaining a carbon balance (see Information Box 16.1).

The following sections discuss the most common organic compounds found in the environment and the microbial catabolic activities that have evolved in response. These include organic polymers, humus and C₁ compounds such as methane (CH₄). It is important to understand the fate of these naturally occurring organic compounds because degradative activities that have evolved for these compounds form the basis for degradation pathways that may be applicable to organic contaminants that are spilled in the environment (see Chapter 31).

Information Box 16.1 The Role of Soil Microbes in Carbon Sequestration

Currently there is debate about how soil microbial activity may influence global warming (Knorr *et al.*, 2005; Rice, 2006). Depending on the relative rates of microbial respiration versus photosynthetic activity, soils could be either a source or sink for CO₂. The estimated amount of carbon sequestered or stored in world soil organic matter ranges from 1.1 to 1.6×10^{12} metric tons (see Table 16.3). This is more than twice the carbon in living vegetation ($\sim 5.6 \times 10^{11}$ metric tons) or in the atmosphere ($\sim 6.7 \times 10^{11}$ metric tons) (Sundquist, 1993). Hence, even relatively small changes in soil carbon storage could have a significant impact on the global carbon balance. In the last 7800 years, the net carbon reservoir in the soil has decreased by 5.0×10^{10} metric tons largely due to conversion of land to agriculture (Lai, 2004). It is estimated that some of this lost carbon could be recovered through strategic management practices. For example, agricultural

practices that enhance crop productivity (CO₂ uptake) while decreasing microbial decomposition rates (CO₂ release) could be optimized to maximize the sequestration of carbon in the soil reservoir. Such practices, which include conservation set-aside, reduced tillage, and increased crop productivity have been estimated to account for the sequestration of 1.1 to 2.1×10^7 metric tons carbon annually (Lokupitva and Paustian, 2006). However, global warming could also result in enhanced rates of microbial decomposition of the carbon stored in the soil. For example, Bellamy *et al.* (2005) documented carbon losses from all soils across England and Wales from the recent period 1978–2003 during which global warming has occurred. Overall, the debate is as yet unresolved, and many subtle feedback effects affect the final outcome, including for example the C:N ratio within soil organic matter and the mandatory C:N requirements of soil microbes.

But before looking more closely at the individual carbon compounds, it should be pointed out that the carbon cycle is actually not quite as simple as depicted in Figure 16.2. This simplified figure does not include anaerobic processes, which were predominant on early Earth and remain important in carbon cycling even today. A more complex carbon cycle containing anaerobic activity is shown in Figure 16.4. Under anaerobic conditions, which predominated for the first few billion years on Earth, some cellular components were less degradable than others (Figure 16.5). This is especially true for highly reduced molecules such as cellular lipids. These components were therefore left over and buried with sediments over time, and became the present-day fossil fuel reserves. Another carbon compound produced under anaerobic conditions is methane. Methane is produced in soils as an end product of anaerobic respiration (see Eq. 3.21, Chapter 3). Methane is also produced in significant quantities under the anaerobic conditions found in ruminants such as cows as well as termite guts. It is also produced within landfills (see Chapter 31).

16.2.3.1 Organic Polymers

What are the predominant types of organic carbon found in the environment? They include plant polymers, fungal and bacterial cell wall polymers, and arthropod exoskeletons (Figure 16.6). Because these polymers constitute the majority of organic carbon, they are the basic food supply available to support heterotrophic activity. The three most common polymers are the plant polymers: cellulose; hemicellulose; and lignin (Table 16.6) (Wagner and Wolf, 1998). The various other polymers produced include: starch (plants); chitin (fungi, arthropods); and




FIGURE 16.4 The carbon cycle, showing both aerobic and anaerobic contributions.




FIGURE 16.5 Examples of petroleum constituents: (A) an alkane; (B) an alicyclic compound; and (C) an aromatic compound. A crude oil contains some of each of these types of compounds but the types and amounts vary in different petroleum reservoirs.




FIGURE 16.6 Common organic polymers found in the environment. (A) Cellulose is the most common plant polymer. It is a linear polymer of β -1,4-linked glucose subunits. Each polymer contains 1000 to 10,000 subunits. (B) Hemicellulose is the second most common polymer. This molecule is more heterogeneous, consisting of hexoses, pentoses and uronic acids. An example of a hemicellulose polymer is pectin. (C) Starch is a polysaccharide synthesized by plants to store energy. Starch is formed from glucose subunits and can be linear (α -1,4 linked), a structure known as amylose, or branched (α -1,4 and α -1,6 linked), known as amylopectin. (D) Chitin is formed from subunits of *N*-acetyl glucosamine linked β -1,4. This polymer is found in fungal cell walls. (E) Bacterial cell walls are composed of polymers of *N*-acetyl glucosamine and *N*-acetylmuramic acid connected by β -1,4 linkages.

peptidoglycan (bacteria). These various polymers can be divided into two groups on the basis of their structures: the carbohydrate-based polymers, which include the majority of the polymers found in the environment, and the phenylpropane-based polymer, lignin.

Carbohydrate-Based Polymers

Cellulose is not only the most abundant of the plant polymers, but it is also the most abundant polymer found on Earth. It is a linear molecule containing β -1,4 linked

glucose subunits (Figure 16.6A). Each molecule contains 1000 to 10,000 subunits with a resulting molecular weight of up to 1.8×10^6 . These linear molecules are arranged in microcrystalline fibers that help make up the woody structure of plants. Cellulose is not only a large molecule; it is also insoluble in water. How then do microbial cells get such a large, insoluble molecule across their walls and membranes? The answer is that they have developed an alternative strategy, which is to synthesize and release enzymes, called **extracellular enzymes**, that can begin the polymer degradation process outside the cell (Deobald

and Crawford, 1997). There are two extracellular enzymes that initiate cellulose degradation. These are β -1,4-endoglucanase and β -1,4-exoglucanase. Endoglucanase hydrolyzes cellulose molecules randomly within the polymer, producing smaller and smaller cellulose molecules (Figure 16.7). Exoglucanase consecutively hydrolyzes two glucose subunits from the reducing end of the cellulose molecule, releasing the disaccharide cellobiose. A third enzyme, known as β -glucosidase or

cellobiase, then hydrolyzes cellobiose to glucose. Cellobiase can be found as both an extracellular and an intracellular enzyme. Both cellobiose and glucose can be taken up by many bacterial and fungal cells.

Hemicellulose is the second most common plant polymer. This molecule is more heterogeneous than cellulose, consisting of a mixture of several monosaccharides including various hexoses and pentoses as well as uronic acids. In addition, the polymer is branched instead of linear. An example of a hemicellulose polymer is the pectin molecule shown in Figure 16.6B, which contains galacturonic acid and methylated galacturonic acid. Degradation of hemicellulose is similar to the process described for cellulose except that, because the molecule is more heterogeneous, many more extracellular enzymes are involved.

In addition to the two major plant polymers, several other important organic polymers are carbohydrate based. One of these is **starch**, a polysaccharide synthesized by plants to store energy (Figure 16.6C). Starch is formed from glucose subunits and can be linear (α -1,4 linked), a structure known as amylose, or can be branched (α -1,4 and α -1,6 linked), a structure known as amylopectin. Amylases

TABLE 16.6 Major Types of Organic Components of Plants

Plant Component	% Dry Mass of Plant
Cellulose	15–60
Hemicellulose	10–30
Lignin	5–30
Protein and nucleic acids	2–15




FIGURE 16.7 The degradation of cellulose begins outside the cell with a series of extracellular enzymes called cellulases. The resulting smaller glucose subunit structures can be taken up by the cell and metabolized.




FIGURE 16.8 (A) Decomposition of wheat straw and its major constituents in a silt loam. The initial composition of the wheat straw was 50% cellulose, 25% hemicellulose and 20% lignin. (B) Leaves from a rain forest in different stages of decomposition. Cellulose and hemicellulose are degraded first, leaving the lignin skeleton. (A) Adapted from [Wagner and Wolf \(1998\)](#). Photo taken in Henry Pittier National Park, Venezuela, courtesy C.M. Miller.

(α -1,4-linked exo- and endoglucanases) are extracellular enzymes produced by many bacteria and fungi. Amylases produce the disaccharide maltose, which can be taken up by cells and mineralized. Another common polymer is **chitin**, which is formed from β -1,4-linked subunits of *N*-acetylglucosamine ([Figure 16.6D](#)). This linear, nitrogen-containing polymer is an important component of fungal cell walls and of the exoskeleton of arthropods. Finally, there is **peptidoglycan**, a polymer of *N*-acetylglucosamine and *N*-acetylmuramic acid, which is an important component of bacterial cell walls ([Figure 16.6E](#)).

Lignin

Lignin is the third most common plant polymer, and is strikingly different in structure from all of the carbohydrate-based polymers. The basic building blocks of lignin are the two aromatic amino acids tyrosine and phenylalanine. These are converted to phenylpropene subunits such as coumaryl alcohol, coniferyl alcohol and sinapyl alcohol. Then 500 to 600 phenylpropene subunits are randomly polymerized, resulting in the formation of the amorphous aromatic polymer known as lignin. In plants, lignin surrounds cellulose microfibrils and strengthens the cell wall. Lignin also helps make plants more resistant to pathogens.

Biodegradation of lignin is slower and less complete than degradation of other organic polymers. This is shown experimentally in [Figure 16.8A](#) and visually in [Figure 16.8B](#). Lignin degrades slowly because it is constructed as a highly heterogeneous polymer, and in addition contains aromatic residues rather than carbohydrate residues. The great heterogeneity of the molecule precludes the evolution of specific degradative enzymes comparable to those for cellulose. Instead, a nonspecific

extracellular enzyme, H_2O_2 -dependent lignin peroxidase, is used in conjunction with an extracellular oxidase enzyme that generates H_2O_2 . The peroxidase enzyme and H_2O_2 system generate oxygen-based free radicals that react with the lignin polymer to release phenylpropene residues ([Morgan et al., 1993](#)). These residues are taken up by microbial cells and degraded as shown in [Figure 16.9](#). Biodegradation of intact lignin polymers occurs only aerobically, which is not surprising because reactive oxygen is needed to release lignin residues. However, once residues are released, they can be degraded under anaerobic conditions.

Phenylpropene residues are aromatic in nature, similar in structure to several types of organic pollutant molecules such as the BTEX (benzene, toluene, ethylbenzene, xylene) and polyaromatic hydrocarbon compounds found in crude oil as well as gasoline and creosote compounds found in wood preservatives (see Chapter 17). These naturally occurring aromatic biodegradation pathways are of considerable importance in the field of bioremediation. In fact, a comparison of the pathway shown in [Figure 16.9](#) with the pathway for degradation of aromatics presented in Chapter 17 shows that they are very similar. Lignin is degraded by a variety of microbes including fungi, actinomycetes and bacteria. The best studied organism with respect to lignin degradation is the white rot fungus *Phanerochaete chrysosporium*. This organism is also capable of degrading several pollutant molecules with structures similar to those of lignin residues (see Information Box 2.6).

16.2.3.2 Humus

Humus was introduced in Chapter 4 and its structure is shown in [Figure 4.14](#). How does humus form? It forms in




FIGURE 16.9 Lignin degradation. Adapted from Wagner and Wolf (1998).

a two-stage process that involves the formation of reactive monomers during the degradation of organic matter, followed by the spontaneous polymerization of some of these monomers into the humus molecule. Although the

majority of organic matter that is released into the environment is respired to form new cell mass and carbon dioxide, a small amount of this carbon becomes available to form humus. To understand this spontaneous process,




FIGURE 16.10 Possible pathways for the formation of soil humus. Adapted with permission from Wagner and Wolf (1998).

consider the degradation of the common organic polymers found in soil, which were described in the preceding sections. Each of these polymers requires the production of extracellular enzymes that begin the polymer degradation process. In particular, for lignin these extracellular enzymes are nonspecific and produce hydrogen peroxide and oxygen radicals. It is not surprising, then, that some of the reactive residues released during polymer

degradation might repolymerize and result in the production of humus. In addition, nucleic acid and protein residues that are released from dying and decaying cells contribute to the pool of molecules available for humus formation. This process is illustrated in Figure 16.10. Considering the wide array of residues that can contribute to humus formation, it is not surprising that humus is even more heterogeneous than lignin. Table 16.7

TABLE 16.7 Chemical Properties of Humus and Lignin

Characteristic	Humic Material	Lignin
Color	Black	Light brown
Methoxyl ($-\text{OCH}_3$) content	Low	High
Nitrogen content	3–6%	0%
Carboxyl and phenolic hydroxyl content	High	Low
Total exchangeable acidity (cmol/kg)	≥ 150	≤ 0.5
α -Amino nitrogen	Present	0
Vanillin content	<1%	15–25%

Data from Wagner and Wolf (1998).

compares the different properties of these two complex molecules.

Humus is the most complex organic molecule found in soil, and as a result it is the most stable organic molecule. The turnover rate for humus ranges from 2 to 5% per year, depending on climatic conditions (Wagner and Wolf, 1998). This can be compared with the degradation of lignin shown in Figure 16.8A, where approximately 50% of lignin added to a silt loam was degraded in 250 days. Thus, humus provides a very slowly released source of carbon and energy for indigenous autochthonous microbial populations. The release of humic residues most likely occurs in a manner similar to release of lignin residues. Because the humus content of most soils does not change, the rate of formation of humus must be similar to the rate of turnover. Thus, humus can be thought of as a molecule that is in a state of dynamic equilibrium (Haider, 1992).

16.2.3.3 Methane

Methanogenesis

The formation of methane, **methanogenesis**, is predominantly a microbial process, although a small amount of methane is generated naturally through volcanic activity (Table 16.8) (Ehrlich, 1996). Methanogenesis is an anaerobic process and occurs extensively in specialized environments including: water-saturated areas such as wetlands and paddy fields; anaerobic niches in the soil; landfills; the rumen; and termite guts. Methane is an end product of anaerobic degradation (see Section 3.4), and as such is associated with petroleum, natural gas and coal deposits.

At present a substantial amount of methane is released to the atmosphere as a result of energy harvesting and

TABLE 16.8 Estimates of Methane Released into the Atmosphere

Source	Methane Emission (10 ⁶ Metric Tons/Year)	
Biogenic		
Ruminants	80–100	
Termites	25–150	
Paddy fields	70–120	
Natural wetlands	120–200	
Landfills	5–70	
Oceans and lakes	1–20	
Tundra	1–5	
Abiogenic		
Coal mining	10–35	
Natural gas flaring and venting	10–35	
Industrial and pipeline losses	15–45	
Biomass burning	10–40	
Methane hydrates	2–4	
Volcanoes	0.5	
Automobiles	0.5	
Total	350–820	
Total biogenic	302–665	81–86% of total
Total abiogenic	48–155	13–19% of total

Adapted from Madigan et al. (1997).


utilization. A second way in which methane is released is through landfill gas emissions (see Chapter 31). Although methane makes a relatively minor carbon contribution to the global carbon cycle (compare Table 16.8 with Table 16.2), methane emission is of concern from several environmental aspects. First, like carbon dioxide, methane is a greenhouse gas and contributes to global warming. In fact, it is the second most common greenhouse gas emitted to the atmosphere. Further, it is 22 times more effective than carbon dioxide at trapping heat. Second, localized production of methane in landfills can create safety and health concerns. Methane is an explosive gas at concentrations as small as 5%. Thus, to avoid accidents, the methane generated in a landfill must be managed in some way. If methane is present in concentrations higher than 35%, it can be collected and used for energy. Alternatively, the methane can be burned off

TABLE 16.9 C₁ Compounds of Major Environmental Importance

Compound	Formula	Comments
Carbon dioxide	CO ₂	Combustion, respiration and fermentation end product, a major reservoir of carbon on Earth
Carbon monoxide	CO	Combustion product, common pollutant. Product of plant, animal and microbial respiration, highly toxic
Methane	CH ₄	End product of anaerobic fermentation or respiration
Methanol	CH ₃ OH	Generated during breakdown of hemicellulose, fermentation by-product
Formaldehyde	HCHO	Combustion product, intermediate metabolite
Formate	HCOOH	Found in plant and animal tissues, fermentation product
Formamide	HCONH ₂	Formed from plant cyanides
Dimethyl ether	CH ₃ OCH ₃	Generated from methane by methanotrophs, industrial pollutant
Cyanide ion	CN ⁻	Generated by plants, fungi and bacteria. Industrial pollutant, highly toxic
Dimethyl sulfide	(CH ₃) ₂ S	Most common organic sulfur compound found in the environment, generated by algae
Dimethyl sulfoxide	(CH ₃) ₂ SO	Generated anaerobically from dimethyl sulfide

at concentrations of 15% or higher. However, most commonly, it is simply vented to the atmosphere to prevent it from building up in high enough concentrations to ignite. Although venting landfill gas to the atmosphere does help prevent explosions, it clearly adds to the global warming problem.


The organisms responsible for methanogenesis are a group of obligately anaerobic archaeobacteria called the **methanogens**. The basic metabolic pathway used by the methanogens is:



This is an exothermic reaction where CO₂ acts as the TEA and H₂ acts as the electron donor providing energy for the fixation of carbon dioxide. Methanogens that utilize CO₂/H₂ are therefore autotrophic. In addition to the autotrophic reaction shown in Eq. 16.1, methanogens can produce methane during heterotrophic growth on a limited number of other C₁ and C₂ substrates including acetate, methanol and formate. Since there are very few carbon compounds that can be used by methanogens, these organisms are dependent on the production of these compounds by other microbes in the surrounding community. As such, an interdependent community of microbes typically develops in anaerobic environments. In this community, the more complex organic molecules are catabolized by populations that ferment or respire anaerobically, generating C₁ and C₂ carbon substrates as well as CO₂ and H₂ that are then used by methanogens.

Methane Oxidation

Clearly, methane as the end product of anaerobiosis is found extensively in nature. As such, it is an available food source, and a group of bacteria called the **methanotrophs** have developed the ability to utilize methane as a source of carbon and energy. The methanotrophs are chemoheterotrophic and obligately aerobic. They metabolize methane as shown in Eq. 16.2:



The first enzyme in the biodegradation pathway is called **methane monooxygenase**. Oxygenases in general incorporate oxygen into a substrate and are important enzymes in the initial degradation steps for hydrocarbons (see Chapter 17). However, methane monooxygenase is of particular interest because it was the first of a series of enzymes isolated that can cometabolize highly chlorinated solvents such as trichloroethene (TCE) (see Chapter 17). Until this discovery it was believed that biodegradation of highly chlorinated solvents could occur only under anaerobic conditions as an incomplete reaction. The application of methanogens for cometabolic degradation of TCE is a strategy under development for bioremediation of groundwater contaminated with TCE. This is a good illustration of the way in which naturally occurring microbial activities can be harnessed to solve pollution problems.





FIGURE 16.11 The nitrogen cycle.


16.2.3.4 Carbon Monoxide and Other C_1 Compounds

Bacteria that can utilize C_1 carbon compounds other than methane are called **methylotrophs**. There are a number of important C_1 compounds produced from both natural and anthropogenic activities (Table 16.9). One of these is carbon monoxide. The annual global production of carbon monoxide is $3\text{--}4 \times 10^9$ metric tons/year (Atlas and Bartha, 1993). The two major carbon monoxide inputs are abiotic. Approximately 1.5×10^9 metric tons/year result from atmospheric photochemical oxidation of carbon compounds such as methane, and 1.6×10^9 metric tons/year results from burning of wood, forests and fossil fuels. A small proportion, 0.2×10^9 metric tons/year, results from biological activity in ocean and soil environments. Carbon monoxide is a highly toxic molecule because it has a strong affinity for cytochromes, and in binding to cytochromes, it can completely inhibit the activity of the respiratory electron transport chain.

Destruction of carbon monoxide can occur abiotically by photochemical reactions in the atmosphere. Microbial processes also contribute significantly to its destruction, even though it is a highly toxic molecule. The destruction of carbon monoxide seems to be quite efficient because the level of carbon monoxide in the atmosphere has not risen significantly since industrialization began, even though CO emissions have increased. The ocean is a net producer of carbon monoxide and releases CO to the atmosphere. In contrast, the terrestrial environment is a net sink for carbon monoxide and absorbs approximately 0.4×10^9 metric tons/year. Key microbes found in terrestrial environments that can metabolize carbon monoxide include both aerobic and anaerobic organisms. Under aerobic conditions *Pseudomonas carboxydoflava* is an example of an organism that oxidizes carbon monoxide to carbon dioxide:



This organism is a chemoautotroph and fixes the CO_2 generated in Eq. 16.3 into organic carbon. The oxidation of the hydrogen produced provides energy for CO_2 fixation (see Eq. 16.4). Under anaerobic conditions, methanogenic bacteria can reduce carbon monoxide to methane:



A number of other C_1 compounds support the growth of methylotrophic bacteria (Table 16.9). Many, but not all, methylotrophs are also methanotrophic. Both types of bacteria are widespread in the environment because these C_1 compounds are ubiquitous metabolites. In response to their presence, microbes have evolved the capacity to metabolize them under either aerobic or anaerobic conditions.

16.3 NITROGEN CYCLE

In contrast to carbon, elements such as nitrogen, sulfur and iron are taken up in the form of mineral salts and cycle oxidoreductively. For example, nitrogen can exist in numerous oxidation states, from -3 in ammonium (NH_4^+) to $+5$ in nitrate (NO_3^-). These element cycles are referred to as the mineral cycles. The best studied and most complex of the mineral cycles is the nitrogen cycle (Figure 16.11). There is great interest in the nitrogen cycle because nitrogen is the mineral nutrient most in demand by microorganisms and plants. It is the fourth most common element found in cells, making up approximately 12% of cell dry weight, and includes the microbially catalyzed processes of nitrogen fixation, ammonium oxidation (aerobic nitrification and anaerobic anammox),

TABLE 16.10 Global Nitrogen Reservoirs

Nitrogen Reservoir	Metric Tons Nitrogen	Actively Cycled
Atmosphere		
N ₂	3.9×10^{15}	No
Ocean		
Biomass	5.2×10^8	Yes
Dissolved and particulate organics	3.0×10^{11}	Yes
Soluble salts (NO ₃ ⁻ , NO ₂ ⁻ , NH ₄ ⁺)	6.9×10^{11}	Yes
Dissolved N ₂	2.0×10^{13}	No
Land		
Biota	2.5×10^{10}	Yes
Organic matter	1.1×10^{11}	Slow
Earth's crust ^a	7.7×10^{14}	No

Adapted from Dobrovolsky (1994).

^aThis reservoir includes the entire lithosphere found in either terrestrial or ocean environments.

assimilatory and dissimilatory nitrate reduction, ammonification and ammonium assimilation. Similarly to the carbon cycle, the global nitrogen cycle is currently undergoing major changes due to the ever-increasing demand for nitrogen in both agriculture and industry, fossil fuel burning and land use changes (Galloway *et al.*, 2008). These perturbations also have major consequences for the environment.

16.3.1 Nitrogen Reservoirs

Nitrogen in the form of the inert gas, dinitrogen (N₂), has accumulated in Earth's atmosphere since the planet was formed. Nitrogen gas is continually released into the atmosphere from volcanic and hydrothermal eruptions, and is one of the major global reservoirs of nitrogen (Table 16.10). A second major reservoir is the nitrogen that is found in Earth's crust as bound, nonexchangeable ammonium. Neither of these reservoirs is actively cycled; the nitrogen in Earth's crust is unavailable, and the N₂ in the atmosphere must be fixed before it is available for biological use. Nitrogen fixation is an energy-intensive and relatively slow process carried out by a limited number of microorganisms. Consequently, the amount of fixed nitrogen available is a limiting factor in primary production, and the nitrogen cycle is closely coupled to the carbon cycle. The pool of fixed nitrogen available can be divided into small reservoirs including the organic nitrogen found in living biomass and in dead organic matter and soluble inorganic nitrogen salts (Table 16.10).

TABLE 16.11 Inputs of Reactive Nitrogen from Natural and Anthropogenic Sources

Source	Nitrogen Fixation (Metric Tons/Year)
Natural nitrogen fixation	
Terrestrial	11×10^7
Aquatic	14×10^7
Lightning	1×10^7
Anthropogenic nitrogen fixation	
Terrestrial (managed farming)	4.6×10^7
Fertilizer manufacture	13.6×10^7
Anthropogenic fixed nitrogen mobilization	
Fossil fuel burning	2.5×10^7
Biomass burning	4×10^7
Wetland draining	1×10^7
Land clearing	2×10^7

These small reservoirs tend to be actively cycled, particularly because nitrogen is often a limiting nutrient in the environment. For example, soluble inorganic nitrogen salts in terrestrial environments can have turnover rates greater than once per day. Nitrogen in plant biomass turns over approximately once a year, and nitrogen in organic matter turns over once in several decades.

16.3.2 Nitrogen Fixation

Ultimately, all fixed forms of nitrogen, NH₄⁺, NO₃⁻ and organic N, come from atmospheric N₂. The relative contributions to **nitrogen fixation** from undisturbed terrestrial and aquatic environments are compared with those influenced by human activity in Table 16.11 (Canfield *et al.*, 2010). From this table one can see that approximately two-thirds of the N₂ fixed annually is microbial, half from terrestrial environments, including both natural systems and managed agricultural systems, and half from marine ecosystems. The remaining third comes from the manufacture of fertilizers. Recall that nitrogen fixation is energy intensive whether microbial or manufactured, and as a result, fertilizer prices are tied to the price of fossil fuels. As fertilizers are expensive, management alternatives to fertilizer addition have become attractive. These include rotation between nitrogen-fixing crops such as soybeans and nonfixing crops such as corn. Wastewater reuse is another alternative that has become especially popular in the desert southwestern United States for nonfood crops and uses, such as cotton and golf courses, where both water and nitrogen are limiting (see Chapter 27).

TABLE 16.12 Representative Genera of Free-Living Nitrogen Fixers

Status with Respect to Oxygen	Mode of Energy Generation	Genus
Aerobic	Heterotrophic	<i>Azotobacter</i> <i>Beijerinckia</i> <i>Acetobacter</i> <i>Pseudomonas</i>
Facultatively anaerobic	Heterotrophic	<i>Klebsiella</i> <i>Bacillus</i>
Microaerophilic	Heterotrophic	<i>Xanthobacter</i> <i>Azospirillum</i>
Strictly anaerobic	Autotrophic Heterotrophic	<i>Thiobacillus</i> <i>Clostridium</i> <i>Desulfovibrio</i>
Aerobic	Phototrophic (cyanobacteria)	<i>Anabaena</i> <i>Nostoc</i>
Facultatively anaerobic	Phototrophic (bacteria)	<i>Rhodospirillum</i>
Strictly anaerobic	Phototrophic (bacteria)	<i>Chlorobium</i> <i>Chromatium</i>

Nitrogen is fixed into ammonia (NH_3) by over 100 different free-living bacteria, both aerobic and anaerobic, as well as some actinomycetes and cyanobacteria (Table 16.12). For example, *Azotobacter* (aerobic), *Beijerinckia* (aerobic), *Azospirillum* (facultative) and *Clostridium* (anaerobic) can all fix N_2 . Because fixed nitrogen is required by all biological organisms, nitrogen-fixing organisms occur in most environmental niches. The amount of N_2 fixed in each niche depends on the environment (Table 16.13). Free-living bacterial cells that are not in the vicinity of a plant root fix small amounts of nitrogen (1 to 2 kg N/hectare/year). Bacterial cells associated with the nutrient-rich rhizosphere environment can fix larger amounts of N_2 (2 to 25 kg N/hectare/year). Cyanobacteria are the predominant N_2 -fixing organisms in aquatic environments, and because they are photosynthetic, N_2 fixation rates are one to two orders of magnitude higher than for free-living nonphotosynthetic bacteria. An evolutionary strategy developed collaboratively by plants and microbes to increase N_2 fixation efficiency was to enter into a symbiotic or mutualistic relationship to maximize N_2 fixation (see Information Box 16.2). The best studied of these symbioses is the *Rhizobium*–legume relationship, which can increase N_2 fixation to 200 to 300 kg N/hectare/year. This symbiosis irrevocably changes both the plant and the microbe involved but is beneficial to both organisms.

TABLE 16.13 Rates of Nitrogen Fixation

N_2 -Fixing System	Nitrogen Fixation (kg N/hectare/year)
<i>Rhizobium</i> –legume	200–300
<i>Anabaena</i> –Azolla	100–120
Cyanobacteria–moss	30–40
Rhizosphere associations	2–25
Free-living	1–2

As the various transformations of nitrogen are discussed in this section, the objective is to understand how they are interconnected and controlled. As already mentioned, N_2 fixation is limited to bacteria and is an energy-intensive process (see Information Box 16.3). Therefore, it does not make sense for a microbe to fix N_2 if sufficient amounts are present for growth. Thus, one control on this part of the nitrogen cycle is that ammonia, the end product of N_2 fixation, is an inhibitor for the N_2 -fixation reaction. A second control in some situations is the presence of oxygen. Nitrogenase is extremely oxygen sensitive, and some free-living aerobic bacteria fix N_2 only at reduced oxygen tension. Other bacteria such as *Azotobacter* and *Beijerinckia* can fix N_2 at normal oxygen tension because they have developed mechanisms to protect the nitrogenase enzyme.

Summary for Nitrogen Fixation

- N_2 fixation is energy intensive
- End product of N_2 fixation is ammonia
- N_2 fixation is inhibited by ammonia
- Nitrogenase is O_2 sensitive; some free-living N_2 fixers require reduced O_2 tension

16.3.3 Ammonia Assimilation (Immobilization) and Ammonification (Mineralization)

The end product of N_2 fixation is ammonia. In the environment, there exists an equilibrium between ammonia (NH_3) and ammonium (NH_4^+) that is driven by pH. This equilibrium favors ammonium formation at acid or near-neutral pH. Thus, it is generally the ammonium form that is assimilated by cells into amino acids to form proteins, cell wall components such as *N*-acetylmuramic acid and purines and pyrimidines to form nucleic acids. This


Information Box 16.2 The Legume–*Rhizobia* Symbiosis

Gram-negative heterotrophic bacteria classified within the genera *Rhizobium*, *Bradyrhizobium*, *Sinorhizobium*, and *Azorhizobium* can form an intriguing symbiosis with leguminous plants. This symbiosis has been well studied because it can meet or nearly meet the nitrogen requirements of economically important legume crops, reducing or eliminating the need for fertilizer application. Grain legumes such as peas, beans, and soybeans can fix about 50% of their total nitrogen requirements, whereas forage legumes such as alfalfa or clover are even more efficient, achieving nitrogen fixation rates as high as 200 to 300 kg per hectare per year. The formation of root nodules on the plant host root system is the result of subtle interactions between the host and the rhizobial endosymbiont (Jones *et al.*, 2007). The plant releases bacteria-specific phenolic compounds called **flavonoids** that in turn signal rhizobia to produce plant-specific lipo-chitooligosaccharide compounds called **Nod factors**. The Nod factors activate a series of host plant responses that prepare the plant to form infection threads, which the invading bacteria use to enter the plant. The infection thread is a thin tubule that penetrates into the plant cortex. Plant cells in the cortex receive the invading bacteria, after which they mature into structures known as symbiosomes. The bacteria in the symbiosome then differentiate into bacteroid cells that are basically nitrogen fixation factories. Also contributing to the success of this amazing invasion process is the fact that the rhizobia can evade all host plant immune responses.

As rhizobia are released from the infection thread, cell division occurs, and a visible root nodule begins to be seen (1 to 2 weeks after infection). Within the root nodule, the rhizobia enlarge and

elongate to perhaps five times the normal size of rhizobia and change physiologically to forms known as bacteroids. The nodule also contains leghemoglobin, which protects the nitrogenase enzyme within the bacteroids from the presence of oxygen. The leghemoglobin imparts a pink color to the interior of the nodule and is indicative of active nitrogen fixation. Thus, examination of the interior of a nodule allows instant determination of whether the nodule is active. Prior to the end of the growing season, nodules begin to break down or senesce, at which point they appear white, green, or brown.

Commercial legume crops are often aided in terms of nitrogen fixation through the application of rhizobial inoculants. This is particularly important when a new legume species is introduced into soils that are free of indigenous rhizobia. In this case, rhizobia introduced into the soil through the use of inoculants tend to establish themselves in the available ecological niche and are difficult to displace by any subsequently introduced rhizobia. Therefore, in such situations it is important that the originally introduced rhizobia are appropriate. The desired characteristics include being infective (capable of causing nodule initiation and development), effective (capable of efficient nitrogen fixation), competitive (capable of causing nodule initiation in the presence of other rhizobia), and persistent (capable of surviving in soil between crops in successive years). Usually rhizobia are impregnated into some kind of peat-based carrier with approximately 10^9 rhizobia per gram of peat. Production of commercial rhizobium inoculants is a big business globally.



The process of legume root hair infection by *Rhizobia*.

process is known as **ammonium assimilation** or **immobilization**. Nitrogen can also be immobilized by the uptake and incorporation of nitrate into organic matter, a process known as **assimilatory nitrate reduction** (Section 16.3.5.1). Because nitrate must be reduced to ammonium before it is incorporated into organic


molecules, most organisms prefer to take up nitrogen as ammonium if it is available. The process that reverses immobilization, the release of ammonia from dead and decaying cells, is called **ammonification** or **ammonium mineralization**. Both immobilization and mineralization of nitrogen occur under aerobic and anaerobic conditions.

Information Box 16.3 The Process of Nitrogen Fixation

Nitrogen gas is a very stable molecule that requires a large amount of energy (946 kJ per mole) to fix into ammonia. The energy for this process arises from the oxidation of carbon sources in the case of heterotrophs or from light in the case of photosynthetic **diazotrophs**. A diazotroph is a microorganism that can fix nitrogen and so does not require fixed nitrogen for growth. Central to biological nitrogen fixation is the enzyme complex **nitrogenase**, which is shown below (Zhao *et al.*, 2006). The overall nitrogenase complex consists of two protein components, which in turn consist of multiple subunits. The iron protein termed **dinitrogenase reductase** is thought to function in the reduction of the molybdenum–iron protein **dinitrogenase**, which reduces nitrogen gas to ammonia. In the 1980s, it was shown that some nitrogenase enzymes do not contain molybdenum and that vanadium and other metals can substitute for molybdenum (Premakumar *et al.*, 1992). To date, three nitrogenase systems have been identified

including Nif, Vnf, and Anf, all of which are sensitive to and inhibited by oxygen. One oxygen-insensitive system has been reported from the actinomycete *Streptomyces thermoautotrophicus*.

The dinitrogenase reductase and dinitrogenase form a complex during which an electron is transferred and the two MgATPs are hydrolyzed to MgADP + inorganic phosphate (Pi). The two proteins then dissociate and the process repeats. After the dinitrogenase protein has collected sufficient electrons, it binds a molecule of nitrogen gas and reduces it, producing ammonia and hydrogen gas. Thus, during reduction of one N₂ molecule, the two proteins must complex and then dissociate a total of eight times. This is the rate-limiting step of the process and takes considerable time. In fact, it takes 1.25 seconds for a molecule of enzyme to reduce one molecule of N₂. This is why nitrogen-fixing bacteria require a great deal of the nitrogenase enzyme, which can constitute 10–40% of the bacterial cell's proteins.



16.3.3.1 Ammonia Assimilation (Immobilization)

There are two pathways that microbes use to assimilate ammonium. The first is a reversible reaction that incorporates or removes ammonium from the amino acid glutamate (Figure 16.12A). This reaction is driven by ammonium availability. At high ammonium concentrations (>0.1 mM or >0.5 mg N/kg soil), in the presence of reducing equivalents (reduced nicotinamide adenine dinucleotide phosphate, NADPH₂), ammonium is

incorporated into α-ketoglutarate to form glutamate using the GOGAT pathway. However, in most soil and many aquatic environments, ammonium is present at low concentrations. Therefore, microbes have a second ammonium uptake pathway that is energy dependent. This reaction is driven by ATP and two enzymes, glutamine synthase and glutamate synthetase (Figure 16.12B). The first step in this reaction adds ammonium to glutamate to form glutamine, and the second step transfers the




FIGURE 16.12 Pathways of ammonium assimilation and ammonification. Assimilation: (A) The enzyme glutamate dehydrogenase catalyzes a reversible reaction that immobilizes ammonium at high ammonium concentrations. (B) The enzyme system glutamine synthetase–glutamate synthetase (GOGAT), which is induced at low ammonium concentrations. This ammonium uptake system requires ATP energy. Ammonification: Ammonium is released from the amino acid glutamate as shown in (A). (C) Ammonia is also released from urea, a molecule that is found in animal waste and in some fertilizers. Note that the first of these reactions (A and B) occurs within the cell. In contrast, urease enzymes are extracellular enzymes, resulting in release of ammonia to the environment. *Source:* Paul and Clark (1989).

ammonium molecule from glutamine to α -ketoglutarate resulting in the formation of two glutamate molecules.

16.3.3.2 Ammonification (Mineralization)

Ammonium mineralization can occur intracellularly by the reversible reaction shown in Figure 16.12A.

Mineralization reactions can also occur extracellularly. Microorganisms release a variety of extracellular enzymes that initiate degradation of plant polymers. Microorganisms also release a variety of enzymes including proteases, lysozymes, nucleases and ureases that can initiate degradation of nitrogen-containing molecules found outside the cell including proteins, cell

walls, nucleic acids and urea. Some of these monomers are taken up by the cell and degraded further, but some are acted upon by extracellular enzymes to release ammonium directly into the environment, as shown in Figure 16.12C for urea and the extracellular enzyme urease.

Which of these two processes, immobilization or mineralization, predominates in the environment? This depends on whether nitrogen is the limiting nutrient. If nitrogen is limiting, then immobilization will become the more important process. For environments where nitrogen is not limiting, mineralization will predominate. Nitrogen limitation is dictated by the carbon/nitrogen (C/N) ratio in the environment. Generally, the C/N ratio required for bacteria is 4 to 5 and for fungi is 10. So, a typical average C/N ratio for soil microbial biomass is 8 (Myrold, 1998). It would then seem logical that at the C/N ratio of 8, there would be a net balance between mineralization and immobilization. However, one must take into account that only approximately 40% of the carbon in organic matter is actually incorporated into cell mass (the rest is lost as carbon dioxide). Thus, the C/N ratio must be increased by a factor of 2.5 to account for the carbon lost as carbon dioxide during respiration. Note that nitrogen is cycled more efficiently than carbon, and there are essentially no losses in its uptake. In fact, a C/N ratio of 20 is not only the theoretical balance point but also the practically observed one. When organic amendments with C/N ratios less than 20 are added to soil, net mineralization of ammonium occurs. In contrast, when organic amendments with C/N ratios greater than 20 are added, net immobilization occurs.

There are numerous possible fates for ammonium that is released into the environment as a result of ammonium mineralization. It can be taken up by plants or microorganisms and incorporated into living biomass, or it can become bound to nonliving organic matter such as soil colloids or humus. In this capacity, ammonium adds to the cation-exchange capacity (CEC) of the soil. Ammonium can become fixed inside clay minerals, which essentially trap the molecule and remove the ammonium from active cycling. Also, because ammonium is volatile, some mineralized ammonium can escape into the atmosphere. Finally, ammonium can be utilized by chemoautotrophic microbes in a process known as nitrification.

Summary for Ammonia Assimilation and Mineralization

- Assimilation and mineralization cycles ammonia between its organic and inorganic forms
- Assimilation predominates at C:N ratios > 20; mineralization predominates at C:N ratios < 20


TABLE 16.14 Chemoautotrophic Nitrifying Bacteria

Genus	Species
Ammonium Oxidizers	
<i>Nitrosomonas</i>	<i>europaea</i> <i>eutrophus</i> <i>marina</i>
<i>Nitrosococcus</i>	<i>nitrosus</i> <i>mobilis</i> <i>oceanus</i>
<i>Nitrospira</i>	<i>briensis</i>
<i>Nitrosolobus</i>	<i>multiformis</i>
<i>Nitrosovibrio</i>	<i>tenuis</i>
Nitrite oxidizers	
<i>Nitrobacter</i>	<i>winogradskyi</i> <i>hamburgensis</i> <i>vulgaris</i>
<i>Nitrospina</i>	<i>gracilis</i>
<i>Nitrococcus</i>	<i>mobilis</i>
<i>Nitrospira</i>	<i>marina</i>

16.3.4 Ammonium Oxidation

16.3.4.1 Nitrification


Nitrification is the microbially catalyzed conversion of ammonium to nitrate (Figure 16.11). This is predominantly an aerobic chemoautotrophic process, but some methylotrophs can use the methane monooxygenase enzyme to oxidize ammonium and a few heterotrophic fungi and bacteria can also perform this oxidation. The autotrophic nitrifiers are a closely related group of bacteria (Table 16.14). The best studied nitrifiers are from the genus *Nitrosomonas*, which oxidizes ammonium to nitrite, and *Nitrobacter*, which oxidizes nitrite to nitrate. The oxidation of ammonium is shown in Eq. 16.6:



This is a two-step energy-producing reaction, and the energy produced is used to fix carbon dioxide. There are two things to note about this reaction. First, it is an inefficient reaction requiring 34 moles of ammonium to fix 1 mole of carbon dioxide. Second, the first step of this reaction is catalyzed by the enzyme ammonium monooxygenase. This first step is analogous to Eq. 16.2, where the

enzyme methane monooxygenase initiates oxidation of methane. Similarly to methane monooxygenase, ammonium monooxygenase has broad substrate specificity, and can be used to oxidize pollutant molecules such as TCE cometabolically (see Section 17.6.2.1).

The second step in nitrification, shown in Eq. 16.7, is even less efficient than the first step, requiring approximately 100 moles of nitrite to fix 1 mole of carbon dioxide:



These two types of nitrifiers, i.e., those that carry out the reactions shown in Eqs. 16.6 and 16.7, are generally found together in the environment. As a result, nitrite does not normally accumulate in the environment. Nitrifiers are sensitive populations. The optimum pH for nitrification is 6.6 to 8.0. In environments with pH < 6.0, nitrification rates are slowed, and below pH 4.5, nitrification seems to be completely inhibited.

Heterotrophic microbes that oxidize ammonium include some fungi and some bacteria. These organisms gain no energy from nitrification, so it is unclear why they carry out the reaction. The relative importance of autotrophic and heterotrophic nitrification in the environment has not yet been clearly determined. Although the measured rates of autotrophic nitrification in the laboratory are an order of magnitude higher than those of heterotrophic nitrification, some data for acidic forest soils have indicated that heterotrophic nitrification may be more important in such environments (Myrold, 1998).


Nitrate does not normally accumulate in natural, undisturbed ecosystems. There are several reasons for this. One is that nitrifiers are sensitive to many environmental stresses. But perhaps the most important reason is that natural ecosystems do not have much excess ammonium. However, in agricultural systems that have large inputs of fertilizer, nitrification can become an important process resulting in the production of large amounts of nitrate. Other examples of managed systems that result in increased nitrogen inputs into the environment are feedlots, septic tanks and landfills. The nitrogen released from these systems also becomes subject to nitrification processes. Because nitrate is an anion (negatively charged), it is very mobile in soil systems, which also have an overall net negative charge. Therefore, nitrate moves easily with water and this results in nitrate leaching into groundwater and surface waters. There are several health concerns related to high levels of nitrate in groundwater, including methemoglobinemia and the formation of nitrosamines. High levels of nitrate in surface waters can also lead to eutrophication and the degradation of surface aquatic systems.

Summary for Nitrification

- Nitrification is an aerobic process that produces nitrite and nitrate
- Nitrification is sensitive to a variety of chemical inhibitors and is inhibited at low pH
- Nitrification in managed systems can result in nitrate leaching and groundwater contamination

16.3.4.2 Anammox

A relative recent discovery is that ammonium oxidation can also occur under anaerobic conditions using nitrite as the terminal electron acceptor (Kuenen, 2008). This process, known as anammox, is the bacterial oxidation of ammonium using nitrite as the electron acceptor resulting in the production of nitrogen (Eq. 16.8):



Anammox was first discovered in a bench-scale wastewater treatment reactor. Since then it has been observed in a wide variety of both aquatic and terrestrial ecosystems with naturally low levels of oxygen. These include marine and freshwater sediments and anoxic water columns, anoxic terrestrial environments, as well as a number of managed systems including wastewater treatment plants, aquaculture and landfill leachate treatment systems (Thamdrup, 2012). All anammox bacteria identified are associated with the phylum Planctomycetes. This phylum is characterized by extremely slow growth rates and in addition has internal membrane-bound structures. In anammox bacteria one such structure is the “anammoxosome,” the organelle where the ammonium oxidation and energy generation reactions take place. Anammox bacteria have not yet been isolated in pure culture (see also Chapter 2).

Anammox bacteria require habitats where both ammonium and nitrite are present. This occurs in the vicinity of aerobic–anaerobic interfaces. At these interfaces, ammonium is released both from the degradation of organic matter (either aerobically or anaerobically) and from dissimilatory nitrate reduction to ammonium (DNRA), an anaerobic process. Nitrite can be produced from nitrate reduction (anaerobic) and also from ammonium that has diffused to an oxic region and then undergone nitrification (Figure 16.11). Thus, competition for anammox substrates occurs at the aerobic–anaerobic interface making this an extremely complex activity.

An immediate and useful application for anammox that was recognized early is the removal of excess ammonium in wastewater treatment. Traditionally, this was done in a two-step process using nitrifying bacteria to

oxidize the ammonium to nitrate under aerobic conditions, and then switching to anoxic conditions to allow denitrification to reduce the nitrate to N_2 (Figure 16.11). This is an energy intensive process and can be replaced by an anammox process. Anammox requires that the first step of nitrification take place (using ammonia oxidizing bacteria) converting some of the ammonium to nitrite (Eq. 16.6). After the first step of nitrification occurs, anammox substrates, nitrite and ammonium are all present. Anammox bacteria can then proceed (at approximately a 1:1 substrate ratio) resulting in the production of N_2 . Research has shown that these steps can take place together in a reactor at one-third to one-half the cost of traditional ammonium removal.

A second discovery was made after anammox was first described. It is now thought that a significant portion (30 to 50%) of the nitrogen loss in ocean environments that had been attributed to denitrification is actually due to anammox. In particular, in ocean zones where oxygen is less than 0.64 mg/L, regions termed oxygen minimum zones, anammox is now thought to dominate as the main cause of fixed nitrogen loss. This is of concern because scientists have suggested that ocean oxygen minimum zones are expanding as a result of climate change. This may in turn cause increased losses of fixed nitrogen through anammox, and a reduction in this very important nutrient in ocean environments.

Summary for Anammox

- Anammox is an anaerobic chemolithoautotrophic process
- Anammox results in the loss of fixed nitrogen to dinitrogen gas
- Anammox is important in ammonium removal in wastewater treatment, and in the loss of fixed nitrogen from oxygen minimum zones in the ocean

16.3.5 Nitrate Reduction

What are the possible fates of nitrate in the environment? Nitrate leaching into groundwater and surface waters is one possible fate. In addition, nitrate can be taken up and incorporated into living biomass by plants and microorganisms. The uptake of nitrate is followed by its reduction to ammonium, which is then incorporated into biomass. This process is called **assimilatory nitrate reduction** or **nitrate immobilization**. Finally, microorganisms can utilize nitrate as a terminal electron acceptor in anaerobic respiration to drive the oxidation of organic compounds. There are two separate pathways for this dissimilatory process, one called **dissimilatory nitrate reduction to ammonium**, where ammonium is the end product, and one called **denitrification**, where a mixture of gaseous products including N_2 and N_2O is formed.

16.3.5.1 Assimilatory Nitrate Reduction

Assimilatory nitrate reduction refers to the uptake of nitrate, its reduction to ammonium and its incorporation into biomass (see Figure 16.12A and B). Most microbes utilize ammonium preferentially, when it is present, to avoid having to reduce nitrate to ammonium, a process requiring energy. So, if ammonium is present in the environment, assimilatory nitrate reduction is suppressed. Oxygen does not inhibit this activity. In contrast to microbes, for plants that are actively photosynthesizing and producing energy, the uptake of nitrate for assimilation is less problematic in terms of energy. In fact, because nitrate is much more mobile than ammonium, it is possible that in the vicinity of the plant roots, nitrification of ammonium to nitrate makes nitrogen more available for plant uptake. Because this process incorporates nitrate into biomass, it is also known as nitrate immobilization (see Section 16.3.3).


Summary for Assimilatory Nitrate Reduction

- Nitrate taken up must be reduced to ammonia before it is assimilated
- Ammonia inhibits this process
- O_2 does not inhibit this process

16.3.5.2 Dissimilatory Nitrate Reduction

Dissimilatory Nitrate Reduction to Ammonium

There are two separate dissimilatory nitrate reduction processes both of which are used by facultative chemoheterotrophic organisms under microaerophilic or anaerobic conditions (Figure 16.11). The first process, called dissimilatory nitrate reduction to ammonium (DNRA), uses nitrate as a terminal electron acceptor to produce energy to drive the oxidation of organic compounds. The end product of DNRA is ammonium:



The first step in this reaction, the reduction of nitrate to nitrite, is the energy-producing step. The further reduction of nitrite to ammonium is catalyzed by an NADH-dependent reductase. This second step provides no additional energy, but it does conserve fixed nitrogen and also regenerates reducing equivalents through the reoxidation of $NADH_2$ to NAD. These reducing equivalents are then used to help in the oxidation of carbon substrates. In fact, it has been demonstrated that under carbon-limiting conditions, nitrite accumulates (denitrification predominates), while under carbon-rich conditions, ammonium is the


major product (DNRA predominates). A second environmental factor that selects for DNRA is low levels of available electron acceptors. It is therefore not surprising that this process is found predominantly in saturated, carbon-rich environments such as stagnant water, sewage sludge, some high-organic-matter sediments and the rumen. Table 16.15 lists a variety of bacteria that perform DNRA. It is interesting to note that most of the bacteria on this list have fermentative rather than oxidative metabolisms.

Summary for Dissimilatory Reduction of Nitrate to Ammonia

- Anaerobic respiration using nitrate as TEA
- Inhibited by O_2
- Not inhibited by ammonia
- Found in a limited number of carbon-rich, TEA-poor environments
- Fermentative bacteria predominate

Denitrification

The second type of dissimilatory nitrate reduction is known as **denitrification**. Denitrification refers to the microbial reduction of nitrate, through various gaseous inorganic forms, to N_2 . This is the primary type of dissimilatory nitrate reduction found in soil, and as such is of concern because it cycles fixed nitrogen back into N_2 . This process removes a limiting nutrient from the environment. Further, some of the gaseous intermediates formed during denitrification, e.g., nitrous oxide (N_2O), can cause depletion of the ozone layer and can also act as a greenhouse gas contributing to global warming (see Chapter 31). The overall reaction for denitrification is:



Denitrification, when calculated in terms of energy produced for every eight-electron transfer, provides more energy per mole of nitrate reduced than DNRA. Thus, in a carbon-limited, electron acceptor-rich environment, denitrification will be the preferred process because it provides more energy than DNRA. The relationship between denitrification and DNRA is summarized in Figure 16.13.

The four steps involved in denitrification are shown in more detail in Figure 16.14. The first step, reduction of nitrate to nitrite, is catalyzed by the enzyme nitrate reductase. This is a membrane-bound molybdenum–iron–sulfur protein that is found not only in denitrifiers but also in

TABLE 16.15 Bacteria that Utilize Dissimilatory Nitrate or Nitrite to Ammonium (DNRA)

Genus	Typical Habitat
Obligate Anaerobes	
<i>Clostridium</i>	Soil, sediment
<i>Desulfovibrio</i>	Sediment
<i>Selenomonas</i>	Rumen
<i>Veillonella</i>	Intestinal tract
<i>Wolinella</i>	Rumen
Facultative Anaerobes	
<i>Citrobacter</i>	Soil, wastewater
<i>Enterobacter</i>	Soil, wastewater
<i>Erwinia</i>	Soil
<i>Escherichia</i>	Soil, wastewater
<i>Klebsiella</i>	Soil, wastewater
<i>Photobacterium</i>	Seawater
<i>Salmonella</i>	Sewage
<i>Serratia</i>	Intestinal tract
<i>Vibrio</i>	Sediment
Microaerophiles	
<i>Campylobacter</i>	Oral cavity
Aerobes	
<i>Bacillus</i>	Soil, food
<i>Neisseria</i>	Mucous membranes
<i>Pseudomonas</i>	Soil, water

Adapted from Tiedje (1988).




FIGURE 16.13 Partitioning of nitrate between denitrification and DNRA as a function of available carbon/electron (C/e^-) acceptor ratio. Adapted from J.M. Tiedje in *Biology of Anaerobic Microorganisms*, A.J.B. Zehnder, ed. © 1988. Reprinted by permission of John Wiley & Sons, Inc.




FIGURE 16.14 The denitrification pathway. Adapted from Myrold (1998).

DNRA organisms. Both the synthesis and the activity of nitrate reductase are inhibited by oxygen. Thus, both denitrification and DNRA are inhibited by oxygen. The second enzyme in this pathway is nitrite reductase, which catalyzes the conversion of nitrite to nitric oxide. Nitrite reductase is unique to denitrifying organisms and is not present in the DNRA process. It is found in the periplasm and exists in two forms, a copper-containing form and a heme form, both of which are distributed widely in the environment. Synthesis of nitrite reductase is inhibited by oxygen and induced by nitrate. Nitric oxide reductase, a membrane-bound protein, is the third enzyme in the pathway, catalyzing the conversion of nitric oxide to nitrous oxide. The synthesis of this enzyme is inhibited by oxygen and induced by various nitrogen oxide forms. Nitrous oxide reductase is the last enzyme in the pathway, and converts nitrous oxide to dinitrogen gas. This is a periplasmic copper-containing protein. The activity of the nitrous oxide reductase enzyme is inhibited by low pH, and is even more sensitive to oxygen than the other three enzymes in the denitrification pathway. Thus, nitrous oxide is the final product of denitrification under conditions of high oxygen (in a relative sense, given a microaerophilic niche) and low pH. In summary, both the synthesis and activity of denitrification enzymes are controlled by oxygen. Enzyme activity is more sensitive to oxygen than enzyme synthesis as

shown in Figure 16.15. The amount of dissolved oxygen in equilibrium with water at 20°C and 1 atm pressure is 9.3 mg/L. However, as little as 0.5 mg/L or less inhibits the activity of denitrification enzymes. Therefore, nitrous oxide reductase is the most sensitive denitrification enzyme, and it is inhibited by dissolved oxygen concentrations of less than 0.2 mg/L.

Whereas the denitrification pathway is very sensitive to oxygen, neither it nor the DNRA pathway is inhibited by ammonium as is the assimilatory nitrate reduction pathway. However, the initial nitrate level in an environmental system can help determine the extent of the denitrification pathway. Low nitrate levels tend to favor production of nitrous oxide as the end product. High nitrate levels favor production of N_2 gas, a much more desirable end product.

Organisms that denitrify are found widely in the environment and display a variety of different characteristics in terms of metabolism and activities. In contrast to DNRA organisms, which are predominantly heterotrophic using fermentative metabolism, the majority of denitrifiers are also heterotrophic, but use respiratory pathways of metabolism. However, as shown in Table 16.16, some denitrifiers are autotrophic, some are fermentative and some are associated with other aspects of the nitrogen cycle; for example, they can fix N_2 .




FIGURE 16.15 Approximate regions of oxygen concentration that inhibit the enzyme activity and synthesis for three steps in the denitrification pathway. Adapted from J.M. Tiedje in *Biology of Anaerobic Microorganisms*, A.J.B. Zehnder, ed. © 1998. Reprinted by permission of John Wiley & Sons, Inc.

Summary for Denitrification

- Anaerobic respiration using nitrate as TEA
- Inhibited by O₂
- Not inhibited by ammonia
- Produces a mix of N₂ and N₂O
- Many heterotrophic bacteria are denitrifiers



16.4 SULFUR CYCLE

Sulfur is the tenth most abundant element in Earth's crust. It is an essential element for biological organisms, making up approximately 1% of the dry weight of a bacterial cell (Table 16.1). Sulfur is not generally considered a limiting nutrient in the environment except in some intensive agricultural systems with high crop yields. Sulfur is cycled between oxidation states of +6 for sulfate (SO₄²⁻) and -2 for sulfide (S²⁻) (Figure 16.16). In cells, sulfur is required for synthesis of the amino acids cysteine and methionine, and is also required for some vitamins, hormones and coenzymes. In proteins, the sulfur-containing amino acid cysteine is especially important because the formation of disulfide bridges between cysteine residues helps govern protein folding, and hence activity. All of these compounds contain sulfur in the reduced or sulfide form. Cells also contain organic sulfur compounds in which the sulfur is in the oxidized state. Examples of such compounds are glucose sulfate, choline sulfate, phenolic sulfate and two ATP-sulfate compounds that are required in sulfate assimilation, and can also serve to store sulfur for the cell. Although the sulfur cycle is not as complex as the nitrogen cycle, the global impacts of the sulfur cycle are extremely

important, including the formation of acid rain, acid mine drainage and corrosion of concrete and metal.

16.4.1 Sulfur Reservoirs

Sulfur is outgassed from Earth's core through volcanic activity. The sulfur gases released, primarily sulfur dioxide (SO₂) and hydrogen sulfide (H₂S), become dissolved in the ocean and aquifers. Here, the hydrogen sulfide forms sparingly soluble metal sulfides, mainly iron sulfide (pyrite), and sulfur dioxide forms metal sulfates with calcium, barium and strontium as shown in Eqs. 16.11 and 16.12:



This results in a substantial portion of the outgassed sulfur being converted to rock. Some of the gaseous sulfur compounds find their way into the upper reaches of the ocean and the soil. In these environments, microbes take up and cycle the sulfur. Finally, the small portions of these gases that remain after precipitating and cycling find their way into the atmosphere. Here, they are oxidized to the water-soluble sulfate form, which is washed out of the atmosphere by rain. Thus, the atmosphere is a relatively small reservoir of sulfur (Table 16.17). Of the sulfur found in the atmosphere, the majority is found as sulfur dioxide. Currently, one-third to one-half of the sulfur dioxide emitted to the atmosphere is from industrial and automobile emissions, due to the burning of fossil fuels. A smaller portion of the sulfur in the atmosphere is present as hydrogen sulfide, and is biological in origin.

TABLE 16.16 Genera of Denitrifying Bacteria

Genus	Interesting Characteristics
Organotrophs	
<i>Alcaligenes</i>	Common soil bacterium
<i>Agrobacterium</i>	Some species are plant pathogens
<i>Aquaspirillum</i>	Some are magnetotactic, oligotrophic
<i>Azospirillum</i>	Associative N ₂ fixer, fermentative
<i>Bacillus</i>	Spore former, fermentative, some species thermophilic
<i>Blastobacter</i>	Budding bacterium, phylogenetically related to <i>Rhizobium</i>
<i>Bradyrhizobium</i>	Symbiotic N ₂ fixer with legumes
<i>Branhamella</i>	Animal pathogen
<i>Chromobacterium</i>	Purple pigmentation
<i>Cytophaga</i>	Gliding bacterium; cellulose degrader
<i>Flavobacterium</i>	Common soil bacterium
<i>Flexibacter</i>	Gliding bacterium
<i>Halobacterium</i>	Halophilic
<i>Hyphomicrobium</i>	Grows on one-C substrates, oligotrophic
<i>Kingella</i>	Animal pathogen
<i>Neisseria</i>	Animal pathogen
<i>Paracoccus</i>	Halophilic, also lithotrophic
<i>Propionibacterium</i>	Fermentative
<i>Pseudomonas</i>	Commonly isolated from soil, very diverse genus
<i>Rhizobium</i>	Symbiotic N ₂ fixer with legumes
<i>Wolinella</i>	Animal pathogen
Phototrophs	
<i>Rhodospseudomonas</i>	Anaerobic, sulfate reducer
Lithotrophs	
<i>Alcaligenes</i>	Uses H ₂ , also heterotrophic, common soil isolate
<i>Bradyrhizobium</i>	Uses H ₂ , also heterotrophic, symbiotic N ₂ fixer with legumes
<i>Nitrosomonas</i>	NH ₃ oxidizer
<i>Paracoccus</i>	Uses H ₂ , also heterotrophic, halophilic
<i>Pseudomonas</i>	Uses H ₂ , also heterotrophic, common soil isolate
<i>Thiobacillus</i>	S-oxidizer
<i>Thiomicrospira</i>	S-oxidizer
<i>Thiosphaera</i>	S-oxidizer, heterotrophic nitrifier, aerobic denitrification

From Myrold (1998).

The largest reservoir of sulfur is found in Earth's crust and is composed of inert elemental sulfur deposits, sulfur–metal precipitates such as pyrite (FeS₂) and gypsum (CaSO₄), as well as sulfur associated with buried fossil fuels. A second large reservoir that is slowly cycled is the sulfate found in the ocean, where it is the second most common anion (Dobrovolsky, 1994). Smaller and more actively cycled reservoirs of sulfur include sulfur found in biomass and organic matter, in the terrestrial and ocean environments. Two recent practices have caused a disturbance in the global sulfur reservoirs. The first is strip mining, which has exposed large areas of metal sulfide ores to the atmosphere, resulting in the formation of acid mine drainage. The second is the burning of fossil fuels, a sulfur reservoir that was quite inert until recently. This has resulted in sulfur dioxide emissions into the atmosphere with the resultant formation of acid rain.

16.4.2 Assimilatory Sulfate Reduction and Sulfur Mineralization

The primary soluble form of inorganic sulfur found in soil is sulfate. Whereas plants and most microorganisms incorporate reduced sulfur (sulfide) into amino acids or other sulfur-requiring molecules, they take up sulfur in the oxidized sulfate form and then reduce it internally (Widdel, 1988). This is called **assimilatory sulfate reduction**. Cells assimilate sulfur in the form of sulfate because it is the most available sulfur form, and also because sulfide is toxic. Sulfide toxicity occurs inside the cell when sulfide reacts with metals in cytochromes to form metal sulfide precipitates, destroying cytochrome activity. However, under the controlled conditions of sulfate reduction inside the cell, the sulfide can be removed immediately and incorporated into an organic form (see **Information Box 16.4**). Although this process does protect the cell from harmful effects of the sulfide, it is an energy-consuming reaction. After sulfate is transported inside the cell, ATP is used to convert the sulfate into the energy-rich molecule adenosine 5'-phosphosulfate (APS) (Eq. 16.14). A second ATP molecule is used to transform APS to 3'-phosphoadenosine-5'-phosphosulfate (PAPS) (Eq. 16.15). This allows the sulfate to be reduced to sulfite and then sulfide in two steps (Eqs. 16.16 and 16.17). Most commonly, the amino acid serine is used to remove sulfide as it is reduced, forming the sulfur-containing amino acid cysteine (see Eq. 16.18).

The release of sulfur from organic forms is called **sulfur mineralization**. The release of sulfur from organic molecules occurs under both aerobic and anaerobic conditions. The enzyme serine sulfhydrylase can remove sulfide from cysteine in the reverse of the reaction shown in Eq. 16.18, or a second enzyme, cysteine sulfhydrylase,









FIGURE 16.16 The sulfur cycle.


TABLE 16.17 Global Sulfur Reservoirs

Sulfur Reservoir	Metric Tons Sulfur	Actively Cycled
Atmosphere		
$\text{SO}_2/\text{H}_2\text{S}$	1.4×10^6	Yes
Ocean		
Biomass	1.5×10^8	Yes
Soluble inorganic ions (primarily SO_4^{2-})	1.2×10^{15}	Slow
Land		
Living biomass	8.5×10^9	Yes
Organic matter	1.6×10^{10}	Yes
Earth's crust ^a	1.8×10^{16}	No


Adapted from Dobrovolsky (1994).

^aThis reservoir includes the entire lithosphere found in either terrestrial or ocean environments.**Information Box 16.4 Processing of Sulfate for Uptake into Bacteria**

can remove both sulfide and ammonia as shown in Eq. 16.19:



In marine environments, one of the major products of algal metabolism is the compound dimethylsulfoniopropionate (DMSP), which is used in osmoregulation of the cell. The major degradation product of DMSP is dimethylsulfide (DMS). Both H_2S and DMS are volatile compounds and therefore can be released to the atmosphere. Once in the atmosphere, these compounds are photooxidized to sulfate (Eq. 16.20):




Normal biological release of reduced volatile sulfur compounds results in the formation of approximately 1 kg SO_4^{2-} /hectare/year. The use of fossil fuels, which all contain organic sulfur compounds, increases the amount of sulfur released to the atmosphere to up to 100 kg SO_4^{2-} /hectare/year in some urban areas. Exacerbating this

TABLE 16.18 Sulfur-Oxidizing Bacteria

Group	Sulfur Conversion	Habitat Requirements	Habitat	Genera
Obligate or facultative chemoautotrophs	$\text{H}_2\text{S} \rightarrow \text{S}^0$ $\text{S}^0 \rightarrow \text{SO}_4^{2-}$ $\text{S}_2\text{O}_3^{2-} \rightarrow \text{SO}_4^{2-}$	H_2S – O_2 interface	Mud, hot springs, mining surfaces, acid mine drainage, soil	<i>Acidithiobacillus</i> , <i>Sulfobacillus</i> , <i>Thiomicrospira</i> , <i>Achromatium</i> , <i>Beggiatoa</i> , <i>Thermothrix</i>
Anaerobic phototrophs	$\text{H}_2\text{S} \rightarrow \text{S}^0$ $\text{S}^0 \rightarrow \text{SO}_4^{2-}$	Anaerobic, H_2S , light	Shallow water, anaerobic sediments meta- or hypolimnion, anaerobic water	<i>Chlorobium</i> , <i>Chromatium</i> , <i>Ectothiorhodospira</i> , <i>Thiopedia</i> , <i>Rhodopseudomonas</i>

Adapted from *Germida (1998)*.

problem is the fact that reserves of fossil fuels that are low in sulfur are shrinking, forcing the use of reserves with higher sulfur content. Burning of fossil fuels produces sulfite as shown in Eq. 16.21:



Thus, increased emission of sulfur compounds to the atmosphere results in the formation of sulfur acid compounds. These acidic compounds dissolve in rainwater and can decrease the rainwater pH from neutral to as low as pH 3.5, a process also known as the formation of **acid rain**. Acid rain damages plant foliage, causes corrosion of stone and concrete building surfaces, and can affect weakly buffered soils and lakes.

Summary for Sulfur Assimilation and Mineralization

- Sulfur is taken up as sulfate rather than sulfide
- Assimilation and mineralization cycles sulfur between its organic and inorganic forms
- Mineralization predominates at C:S ratio <200; assimilation predominates at C:S >400


16.4.3 Sulfur Oxidation

In the presence of oxygen, reduced sulfur compounds can support the growth of a group of chemoautotrophic bacteria under strictly aerobic conditions and a group of photoautotrophic bacteria under strictly anaerobic conditions (Table 16.18). In addition, a number of aerobic heterotrophic microbes, including both bacteria and fungi, oxidize sulfur to thiosulfate or to sulfate. The heterotrophic sulfur oxidation pathway is still unclear, but apparently no energy is obtained in this process. Chemoautotrophs are considered the predominant sulfur oxidizers in most environments. However, because many

chemoautotrophic sulfur oxidizers require a low pH for optimal activity, heterotrophs may be more important in some aerobic, neutral to alkaline soils. Further, heterotrophs may initiate sulfur oxidation, resulting in a lowered pH that is more amenable for chemoautotrophic activity.


16.4.3.1 Chemoautotrophic Sulfur Oxidation

Of the chemoautotrophs, most oxidize sulfide to elemental sulfur, which is then deposited inside the cell as characteristic granules (Eq. 16.22):



The energy provided by this oxidation is used to fix CO_2 for cell growth. In examining Eq. 16.22, it is apparent that these organisms require both oxygen and sulfide. However, reduced compounds are generally abundant in areas that contain little or no oxygen. So these microbes are **microaerophilic**; they grow best under conditions of low oxygen tension (Figure 16.17). Characteristics of marsh sediments that contain these organisms are their black color due to sulfur deposits, and their “rotten egg” smell due to the presence of H_2S . Most of these organisms are filamentous and can easily be observed by examining a marsh sediment under the microscope and looking for small white filaments.

Some chemoautotrophs, most notably *Acidithiobacillus thiooxidans*, can oxidize elemental sulfur as shown in Eq. 16.23:



This reaction produces acid, and as a result, *A. thiooxidans* is extremely acid tolerant with an optimal growth pH of 2. It should be noted that there are various *Acidithiobacillus* species, and these vary widely in




FIGURE 16.17 Cultivation of the sulfur oxidizing chemolithotroph *Beggiatoa*. At right is a culture tube with sulfide agar overlaid with initially sulfide-free soft mineral agar. The air space in the closed tube is the source of oxygen. Stab-inoculated *Beggiatoa* grows in a narrowly defined gradient of H_2S and oxygen as shown. Adapted with permission from *Microbial Ecology* by R.M. Atlas and R. Bartha. © 1993, by Benjamin Cummings.

their acid tolerance (Baker and Banfield, 2003). However, the activity of *A. thiooxidans* in conjunction with the iron-oxidizing, acid-tolerant, chemoautotroph *Acidithiobacillus ferrooxidans* is responsible for the formation of acid mine drainage, an undesirable consequence of sulfur cycle activity. It should be noted that the same organisms can be harnessed for the acid leaching and recovery of precious metals from low-grade ore, also known as **biometallurgy**. Thus, depending on one's perspective, these organisms can be very harmful or very helpful.

Although most of the sulfur-oxidizing chemoautotrophs are obligate aerobes, there is one exception, *Acidithiobacillus denitrificans*, a facultative anaerobic organism that can substitute nitrate as a terminal electron acceptor for oxygen as shown in Eq. 16.24:




In the above equation, the sulfate formed is shown as precipitating with calcium to form gypsum. *A. denitrificans* is not acid tolerant, and has an optimal pH for growth of 7.0.

Summary for Chemoautotrophic Sulfur Oxidation

- It is an aerobic process, but most sulfur oxidizers are microaerophilic
- Some heterotrophs oxidize sulfur but obtain no energy
- This process can result in the formation of acid mine drainage
- This process can be used in metal recovery

16.4.3.2 Photoautotrophic Sulfur Oxidation

Photoautotrophic oxidation of sulfur is limited to green and purple sulfur bacteria (Table 16.17). This group of bacteria evolved on early Earth when the atmosphere contained no oxygen. These microbes fix carbon using light energy, but instead of oxidizing water to oxygen, they use an analogous oxidation of sulfide to sulfur:



These organisms are found in mud and stagnant water, sulfur springs and saline lakes. In each of these environments, both sulfide and light must be present. Although the contribution to primary productivity is small in comparison with aerobic photosynthesis, these organisms are important in the sulfur cycle. They serve to remove sulfide from the surrounding environment, effectively preventing its movement into the atmosphere and its precipitation as metal sulfide.


Summary for Photoautotrophic Sulfur Oxidation

- It is an anaerobic process that is limited to the purple and green sulfur bacteria
- It is responsible for a very small portion of total photosynthetic activity


16.4.4 Sulfur Reduction

There are three types of sulfur reduction. The first, already discussed in Section 16.4.2, is performed to assimilate sulfur into cell components (Widdel, 1988). Assimilatory sulfate reduction occurs under either aerobic or anaerobic


conditions. In contrast, there are two dissimilatory pathways, both of which use an inorganic form of sulfur as a terminal electron acceptor. In this case, sulfur reduction occurs only under anaerobic conditions. The two types of sulfur that can be used as terminal electron acceptors are elemental sulfur and sulfate. These two types of metabolism are differentiated as **sulfur respiration** and **dissimilatory sulfate reduction**. *Desulfuromonas acetooxidans* is an example of a bacterium that grows on small carbon compounds such as acetate, ethanol and propanol, and uses elemental sulfur as the terminal electron acceptor as shown in Eq. 16.26:



However, the use of sulfate as a terminal electron acceptor seems to be the more important environmental process. The following genera, all of which utilize sulfate as a terminal electron acceptor, are found widely distributed in the environment, especially in anaerobic sediments of aquatic environments, water-saturated soils and animal intestines: *Desulfobacter*, *Desulfobulbus*, *Desulfococcus*, *Desulfonema*, *Desulfosarcina*, *Desulfotomaculum* and *Desulfovibrio*. Together these organisms are known as the **sulfate-reducing bacteria (SRB)**. They can utilize H_2 as an electron donor to drive the reduction of sulfate as shown in Eq. 16.27:



Thus, SRB compete for available H_2 in the environment, as H_2 is also the electron donor required by methanogens. It should be noted that this is not usually a chemoautotrophic process because most SRB cannot fix carbon dioxide. Instead, they obtain carbon from low-molecular-weight compounds such as acetate or methanol. The overall reaction for utilization of methanol is shown in Eq. 16.28:



Both sulfur and sulfate reducers are strict anaerobic chemoheterotrophic organisms that prefer small carbon substrates such as acetate, lactate, pyruvate and low-molecular-weight alcohols. Where do these small carbon compounds come from in the environment? They are by-products of fermentation of plant and microbial biomass that occurs in anaerobic regions. Thus, the sulfate reducers are part of an anaerobic consortium of bacteria including fermenters, sulfate reducers and methanogens, which together act to completely mineralize organic compounds to carbon dioxide and methane (see Chapter 3). More recently, it has been found that some SRB can also metabolize more complex carbon compounds including some aromatic compounds and some longer chain fatty

acids. These organisms are being looked at closely to determine whether they can be used in remediation of contaminated sites that are highly anaerobic, and that would be difficult to oxygenate.

The end product of sulfate reduction is hydrogen sulfide. What are the fates of this compound? It can be taken up by chemoautotrophs or photoautotrophs and reoxidized; it can be volatilized into the atmosphere; or it can react with metals to form metal sulfides. In fact, the activity of sulfate reducers and the production of hydrogen sulfide are responsible for the corrosion of underground metal pipes. In this process, the hydrogen sulfide produced reacts with ferrous iron metal to make more iron sulfide.

Summary for Sulfur Reduction

- Anaerobic respiration using SO_4^{2-} or S^0 as a TEA
- Completely inhibited by O_2
- Produces H_2S which can cause metal corrosion

16.5 IRON CYCLE

16.5.1 Iron Reservoirs

Iron is the fourth most abundant element in Earth's crust. Iron generally exists in three oxidation states: 0, +2 and +3 corresponding to metallic iron (Fe^0), ferrous iron (Fe^{2+}) and ferric iron (Fe^{3+}). In the environment, iron is actively cycled between the +2 and +3 forms (Figure 16.18). Under aerobic conditions iron is usually found in its most oxidized form (Fe^{3+}), which has low aqueous solubility. Under reducing or anaerobic conditions Fe^{3+} is reduced to the ferrous form, Fe^{2+} , which has higher solubility. Iron is an essential but minor element for biological organisms, making up approximately 0.2% of the dry weight of a bacterial cell (Table 16.1). Although the amount of iron in a cell is low, it has a very important function as a part of enzymes that are used in respiration and photosynthesis, both processes that require electron transfer.

16.5.2 Iron in Soils and Sediments

Iron is generally not a limiting nutrient in soil due to its high abundance in Earth's crust. However, even though iron abundance is high, the bioavailability of most iron minerals is quite limited. Thus, microorganisms have developed strategies to obtain iron from its mineral form, usually from iron oxides or iron oxyhydroxides. The best studied strategy is the use of iron chelators known as siderophores (Figure 16.19). Siderophores are synthesized and released from the cell, where they bind Fe^{3+} , which helps keep this low solubility form of iron in solution.




FIGURE 16.18 Conceptual diagram of iron redox cycling in soils and sediments. Iron atoms can cycle many times through oxidized and reduced states before being lost from the soil profile. Consequently, electron fluxes associated with iron may greatly exceed the oxidation capacity represented by the mass of iron oxides present at any given time. Courtesy Aaron Thompson.




FIGURE 16.19 Two siderophores showing the coordination of the siderophores with iron. Ferrichrome is produced by fungi including *Aspergillus*, *Ustilago* and *Penicillium*. Pseudobactin is a bacterial siderophore made by *Pseudomonas* sp.

The soluble iron–siderophore complex is recognized by and binds to siderophore-specific receptors on the cell surface. The iron is released from the complex, reduced and taken up into the cell as Fe^{2+} , its more soluble form.

16.5.3 Iron in Marine Environments

Unlike terrestrial and sediment environments, iron is considered a limiting nutrient in the modern marine environment (Raiswell, 2006). In fact, the extent of this limitation has been the focus of a vigorous debate in recent years. Recent data seem to indicate that for one-third of Earth's oceans, those that are more nutrient rich and support higher numbers of phytoplankton, iron is the limiting nutrient to growth (Boyd *et al.*, 2007). How does iron enter the ocean environment? As shown in Table 16.19, the largest flux of iron entering the oceans is

fluvial in nature, meaning that it enters as suspended particles or dissolved iron carried by rivers, or is associated with glacial sediments (Jickells *et al.*, 2005). For the most part, this iron is deposited in the sediments of near-coastal areas, and does not reach the open ocean. Therefore, in the open ocean, the major pathway of iron entry is *colian*, or as dust that is carried through the atmosphere mainly from desert and other arid environment land surfaces.

16.5.4 Iron Oxidation

16.5.4.1 Chemoautotrophs


Under aerobic conditions, ferrous iron tends to oxidize to the ferric form. Ferrous iron will autoxidize or spontaneously oxidize under aerobic conditions at $\text{pH} > 5$. Iron

TABLE 16.19 Global Iron Fluxes to the Ocean

Source	Flux Teragrams (1×10^9 kg) Per Year
Fluvial particulate total iron	625–962
Fluvial dissolved iron	1.5
Glacial sediments	34–211
Atmospheric	16
Coastal erosion	8
Hydrothermal	14
Authigenic (release from deep-sea sediments during diagenesis)	5

Adapted from Jickells et al. (2005).

oxidation also occurs biotically. In fact, reduced iron is an important source of energy for several specialized genera of chemoautotrophic bacteria, the iron-oxidizers (Ehrlich, 1996):



Note that compared to ammonia (Eq. 16.6) or sulfur (Eqs. 16.22, 16.23) oxidation, the yield of energy in iron oxidation is quite low. Nevertheless, this is an exploitable niche and an important biological reaction, which can have considerable environmental consequences.

Iron oxidation is most often associated with acidic environments. For example, the acidophilic thermophile *Sulfolobus* is an archaeon that was isolated from acidic hot springs. But iron oxidation is perhaps best studied in association with acid mine drainage and the acidophilic bacterium *Acidithiobacillus ferrooxidans*. Interestingly, although *A. ferrooxidans* is best studied (because it has been most easily cultured), nonculture-based analysis suggests that other iron-oxidizers may actually play more important roles in the creation of acid conditions in mine tailings and the formation of acid mine drainage (Baker and Banfield, 2003). These include Bacteria such as *Leptospirillum ferrooxidans* and *L. ferriphilum*, *Sulfolobus acidophilus*, *S. thermosulfophilus* and *Acidimicrobium ferrooxidans*, as well as some Archaea, e.g., *Ferroplasma acidiphilum*.

Iron oxidation has also been described in several marine genera at neutral pH. This process is problematic at neutral pH because: (1) the energy yield is very low; and (2) at neutral pH, spontaneous oxidation of Fe^{2+} to Fe^{3+} (in the form of iron oxyhydroxides) occurs rapidly in the presence of oxygen and competes with the




FIGURE 16.20 Iron hydroxide- and iron oxyhydroxide-coated stalks of *Gallionella* and sheaths of *Leptothrix*. In this transmission electron micrograph, dark areas are the metal deposits. From University of California Berkeley Geomicrobiology Program, 2008.

biological reaction (Figure 16.18). Neutrophilic iron-oxidizers overcome these problems with very specific niche requirements. They position themselves in regions with low O_2 tension and high and constant Fe^{2+} concentrations. One marine environment that meets these requirements occurs in hydrothermal vents on the sea floor (see Chapter 7). Here, anoxic vent fluids charged with Fe^{2+} rapidly come in contact with the cold, oxygenated ocean water. Similar conditions can occur in municipal and industrial water pipelines.


The best-studied neutrophilic iron-oxidizers are *Gallionella* and *Leptothrix*. *Gallionella* is capable of chemoautotrophic growth using Fe^{2+} as sole energy source under microaerobic conditions. *Leptothrix* is a sheath-forming chemoheterotrophic organism that oxidizes both Fe^{2+} and Mn^{2+} , depositing an iron–manganese encrusted coating on its sheath. As can be seen in Figure 16.20, these microbes can deposit copious amounts of iron (and manganese) minerals on their surfaces. This can have serious economic consequences in some instances. For example, this process can cause extensive biofouling and corrosion of water pipelines.

Summary for Iron Oxidation

- Iron oxidation is a chemoautotrophic, aerobic process usually found under acidic conditions
- This process participates in the formation of acid mine drainage (and can be used for metal recovery)
- Neutrophilic iron oxidation occurs in a more limited number of environments where, due to iron mineral deposition, this process can result in biofouling and corrosion

16.5.4.2 Photoautotrophs

As shown below, some members of the purple and green bacteria can use Fe^{2+} as an electron donor to carry out anaerobic photosynthesis coupled to photoautotrophic growth (i.e., growth involving photosynthesis and oxidation):



It has been proposed that iron-based photosynthesis represents the transition between anaerobic photosynthesis that developed on early Earth and aerobic photosynthesis that began approximately 2 billion years ago (Jiao and Newman, 2007). In fact, it is thought that Fe^{2+} was the most widespread source of reducing power from 1.6 to 3.8 billion years ago, where under reducing conditions, Fe^{2+} was favored over Fe^{3+} . Under these conditions, the amount of iron in the marine environment was considerably higher (0.1 to 1 millimoles/L) than it is today (0.03 to 1 nanomoles/L) (Jickells *et al.*, 2005; Bosak *et al.*, 2007).

Summary for Photoautotrophs

- Phototrophic iron oxidation is a photoautotrophic, anaerobic process that is limited to purple and green sulfur bacteria
- This process is thought to be a key transition step between anaerobic photosynthesis found on early Earth and modern aerobic photosynthesis

16.5.5 Iron Reduction

Iron is microbially reduced for two purposes, assimilation and energy generation. **Assimilatory iron reduction** is the reduction of Fe^{3+} for uptake and incorporation into cell constituents. As described in Section 16.5.2, this usually involves the release of siderophores, which complex Fe^{3+} in the environment exterior to the cell. The iron–siderophore complex then delivers the Fe^{3+} to the cell, which is reduced to Fe^{2+} as it is taken up.

Dissimilatory iron reduction or **iron respiration** is the use of Fe^{3+} as a terminal electron acceptor for the purpose of energy generation during anaerobic respiration. Due to its abundance in Earth's crust, Fe^{3+} found in iron oxides and oxyhydroxides serves as an important terminal electron acceptor for anaerobic heterotrophic bacteria. Because iron respiration has been an important activity during the evolution of Earth, there is wide diversity among the bacteria and archaeans capable of carrying out this activity (Weber *et al.*, 2006). The problem for the iron-reducers is that most of the iron in environment is relatively unavailable (Figure 16.18). So, microorganisms have developed some very interesting strategies to solve this problem (Lovley, 2000). As shown in Figure 16.21,




FIGURE 16.21 Two primary strategies are used in iron respiration to facilitate electron transfer between microbial cells and iron oxide surfaces. (A) Cells can come in direct contact with the oxide surface. (B) An electron shuttle can be used to mediate the electron transfer between the cell and iron oxide surfaces.

the first is to make direct contact with an iron oxide surface. In this case, the iron reductase is a membrane-bound enzyme allowing direct access of the enzyme with the substrate. A second strategy is to use an electron shuttle that can act as an intermediate in transferring electrons from the cell to the iron oxide surface. Possible electron shuttles in the environment include humic acids or other molecules that contain quinone-like structures that are reduced to a corresponding hydroquinone form. Humic acids are considered an **exogenous** electron shuttle source, or one that is obtained from the environment outside the cell. Alternatively, some bacteria can make and release **endogenous** electron shuttles. For example, *Geothrix fermentans* releases a quinone-like electron shuttle during growth on lactate (electron donor) and poorly crystalline iron oxides (electron acceptor) (Nevin and Lovley, 2002).

Summary for Iron Reduction

- Assimilatory iron reduction is generally mediated by iron-chelating molecules called siderophores
- Dissimilatory iron reduction is the use of ferric iron as a terminal electron acceptor during anaerobic respiration. Due to the abundance of iron in soils and sediments, it is thought that this is an important process in anaerobic environments

QUESTIONS AND PROBLEMS

1. Describe both microbial and non-microbial contributions to the carbon cycle.
2. Give an example of
 - a. a small actively cycled reservoir
 - b. a large actively cycled reservoir
 - c. a large inactively cycled reservoir

3. Describe how the ocean has reduced the expected rate of increase of CO₂ in the atmosphere since industrialization began.
4. What is the concept behind fertilization of the ocean with iron?
5. What strategy is used by microbes to initiate degradation of large plant polymers such as cellulose?
6. Define what is meant by a greenhouse gas and give two examples. For each example describe how microorganisms mediate generation of the gas, and then describe how human activity influences generation of the gas.
7. Both autotrophic and heterotrophic activities are important in element cycling. For each cycle discussed in this chapter (carbon, nitrogen, sulfur and iron), name the most important heterotrophic and autotrophic activity. Justify your answer.
8. What would happen if microbial nitrogen fixation suddenly ceased?

REFERENCES AND RECOMMENDED READING

- Atlas, R. M., and Bartha, R. (1993) "Microbial Ecology," Benjamin Cummings, New York.
- Baker, B. J., and Banfield, J. F. (2003) Microbial communities in acid mine drainage. *FEMS Microbiol. Ecol.* **44**, 139–152.
- Bellamy, P. H., Loveland, P. J., Bradley, R. I., Lark, R. M., and Kirk, G. J. (2005) Carbon losses from all soils across England and Wales 1978–2003. *Nature* **437**, 245–248.
- Bosak, T., Greene, S. E., and Newman, D. K. (2007) A likely role for anoxygenic photosynthetic microbes in the formation of ancient stromatolites. *Geobiology* **5**, 119–126.
- Boyd, P. W., Jickells, T., Law, C. S., Blain, S., Boyle, E. A., Buesseler, K. O., *et al.* (2007) Mesoscale iron enrichment experiments 1993–2005: synthesis and future directions. *Science* **315**, 612–617.
- Canfield, D. E., Glazer, A. N., and Falkowski, P. G. (2010) The evolution and future of earth's nitrogen cycle. *Science* **330**, 192–196.
- Deobald, L. A., and Crawford, D. L. (1997) Lignocellulose biodegradation. In "Manual of Environmental Microbiology" (C. J. Hurst, G. R. Knudsen, M. J. McInerney, L. D. Stetzenbach, and M. V. Walter, eds.), American Society for Microbiology, Washington, DC, pp. 730–737.
- Dobrovolsky, V. V. (1994) "Biogeochemistry of the World's Land," CRC Press, Boca Raton, FL.
- Ehrlich, H. L. (1996) "Geomicrobiology," Marcel Dekker, New York.
- Farabee, M. (2008) Paleobiology: the precambrian: life's genesis and spread. In "Online Biology Book", Estrella Mountain Community College, Avondale, Arizona. <http://www.emc.maricopa.edu/faculty/farabee/BIOBK/BioBookPaleo2.html>.
- Galloway, J. N., Townsend, A. R., Erisman, J. W., Bekunda, M., Cai, Z., Feney, J. R., *et al.* (2008) Transformation of the nitrogen cycle: recent trends, questions, and potential solutions. *Science* **320**, 889–892.
- Germida, J. J. (1998) Transformations of sulfur. In "Principles and Applications of Soil Microbiology" (D. M. Sylvia, J. J. Fuhrmann, P. G. Hartel, and D. A. Zuberer, eds.), Prentice-Hall, Upper Saddle River, NJ, pp. 346–368.
- Haider, K. (1992) Problems related to the humification processes in soils of temperate climates. In "Soil Biochemistry" (G. Stotzky, and J.-M. Bollag, eds.), vol. 7, Marcel Dekker, New York, pp. 55–94.
- Jiao, Y., and Newman, D. K. (2007) The pio operon is essential for phototrophic Fe(II) oxidation in *Rhodopseudomonas palustris* TIE-1. *J. Bacteriol.* **189**, 1765–1773.
- Jickells, T. D., An, Z. S., Andersen, K. K., Baker, A. R., Bergametti, G., Brooks, N., *et al.* (2005) Global iron connections between desert dust, ocean biogeochemistry, and climate. *Science* **308**, 67–71.
- Jones, K. M., Kobayashi, H., Davies, B. W., Taga, M. E., and Walker, G. C. (2007) How rhizobial symbionts invade plants: the *Sinorhizobium-Medicago* model. *Nat. Rev. Microbiol.* **5**, 619–633.
- Knorr, W., Prentice, I. C., House, J. I., and Holland, E. A. (2005) Long-term sensitivity of soil carbon turnover to global warming. *Nature* **433**, 298–301.
- Kuenen, J. G. (2008) Anammox bacteria: from discovery to application. *Nat. Rev. Microbiol.* **6**, 320–326.
- Lai, R. (2004) Soil carbon sequestration impacts on global climate change and food security. *Science* **304**, 1623–1627.
- Lokupitiya, E., and Paustian, K. (2006) Agricultural soil greenhouse gas emissions: a review of national inventory methods. *J. Environ. Qual.* **35**, 1413–1427.
- Lovelock, J. (1995) "The Ages of Gaia," W. W. Norton, New York.
- Lovley, D. R. (2000) Fe(III) and Mn(IV) reduction. In "Environmental Microbe–Metal Interactions" (D. R. Lovley, ed.), ASM Press, Washington, DC, pp. 3–30.
- Madigan, M. T., Martinko, J. M., and Parker, J. (1997) "Brock Biology of Microorganisms," 8th ed. Prentice Hall, Upper Saddle River, NJ.
- Morgan, P., Lee, S. A., Lewis, S. T., Sheppard, A. N., and Watkinson, R. J. (1993) Growth and biodegradation by white-rot fungi inoculated into soil. *Soil Biol. Biochem.* **25**, 279–287.
- Myrold, D. D. (1998) Transformations of nitrogen. In "Principles and Applications of Soil Microbiology" (D. M. Sylvia, J. J. Fuhrmann, P. G. Hartel, and D. A. Zuberer, eds.), Prentice-Hall, Upper Saddle River, NJ, pp. 218–258.
- Neidhardt, F. C., Ingraham, J. L., and Schaechter, M. (1990) "Physiology of the Bacterial Cell: A Molecular Approach," Sinauer Associates, Sunderland, MA.
- Nevin, K. P., and Lovley, D. R. (2002) Mechanisms for accessing insoluble Fe(III) oxide during dissimilatory Fe(III) reduction by *Geothrix fermentans*. *Appl. Environ. Microbiol.* **68**, 2294–2299.
- Paul, E. A., and Clark, F. E. (1989) "Soil Microbiology and Biochemistry," Academic Press, New York.
- Premakumar, R., Jacobson, M. R., Loveless, T. M., and Bishop, P. E. (1992) Characterization of transcripts expressed from nitrogenase-3 structural genes of *Azotobacter vinelandii*. *Can J Microbiol.* **38**, 929–936.
- Raiswell, R. (2006) Towards a global highly reactive iron cycle. *J. Geochem. Explor.* **88**, 436–439.
- Reynolds, S. (1999) Stromatolites. In "MLSSA Journal," The Marine Life Society of South Australia, Inc., Adelaide, Australia, <http://www.mlssa.asn.au/>.
- Rice, C. W. (2006) Introduction to special section on greenhouse gases and carbon sequestration in agriculture and forest. *J. Environ. Qual.* **35**, 1338–1340.

- Sundquist, E. T. (1993) The global carbon dioxide budget. *Science* **259**, 934–941.
- Thamdrup, B. (2012) New pathways and processes in the global nitrogen cycle. *Annu. Rev. Col. Evol. Syst.* **43**, 407–428.
- Tiedje, J. M. (1988) Ecology of denitrification and dissimilatory nitrate reduction to ammonium. In “Biology of Anaerobic Microorganisms” (A. J. B. Zehnder, ed.), John Wiley & Sons, New York, pp. 179–244.
- Wagner, G. H., and Wolf, D. C. (1998) Carbon transformations and soil organic matter formation. In “Principles and Applications of Soil Microbiology” (D. M. Sylvia, J. J. Fuhrmann, P. G. Hartel, and D. A. Zuberer, eds.), Prentice-Hall, Upper Saddle River, NJ, pp. 259–294.
- Weber, K. A., Achenbach, L. A., and Coates, J. D. (2006) Microorganisms pumping iron: anaerobic microbial iron oxidation and reduction. *Nat. Rev. Microbiol.* **4**, 752–764.
- Widdel, F. (1988) Microbiology and ecology of sulfate- and sulfur-reducing bacteria. In “Biology of Anaerobic Microorganisms” (A. J. B. Zehnder, ed.), John Wiley & Sons, New York, pp. 469–585.
- Zhao, Y., Bian, S. M., Zhou, H. N., and Huang, J. F. (2006) Diversity of nitrogenase systems in diazotrophs. *J. Integr. Plant Biol.* **48**, 745–755.